

100%
Success

Mathematics

Only
40 Days

For

Class

11th

- ❖ اہم ترین سوالات کے جوابات
- ❖ تجربہ کار اساتذہ کی زیر نگرانی میں
- ❖ نوٹس، گیس پیپر، ٹیسٹ سیریز

- ❖ کم وقت میں بہترین تیاری
- ❖ پیپر پیٹرن کے ذہن کو مد نظر رکھتے ہوئے
- ❖ مختصر وقت میں نوٹس کے ذریعے شاندار تیاری

❖ اب پریشان ہونے کی ضرورت نہیں پاک سٹی (pakcity.org) آپ کے لئے لایا ہے بہترین شارٹ نوٹس اور گیس پیپر۔

❖ پاک سٹی (pakcity.org) کے ایسے نوٹس اور گیس پیپر جن کو کمزور طلبہ و طلبات 40 دن کی تیاری کر کے پڑھائی میں +A بھی حاصل کر سکتے ہیں۔

❖ پاک سٹی (pakcity.org) کے نوٹس میں بہت اہم معروضی سوالات، مختلف سوالات اور انشائیہ سوالات سلیوشن کے ساتھ شامل ہیں۔

نوٹ: اگر آپ کو کسی بھی بورڈ کی کوئی بھی کتاب کے نوٹس چاہیے تو آپ pakcity.org وزٹ کریں۔

**Bahawalpur Board
Gujranwala Board
Rawalpindi Board**

**D.G.Khan Board
Faisalabad Board
Sargodha Board**

**Lahore Board
Multan Board
Sahiwal board**

Notes, Guess Papers, Test Series of the All Classes:

5th6th7th8th9th10th11th12th

Bachelor and Masters Programs in all departments andetc

English
Urdu
Physics
Farsi

Islamiat
Biology
Civics
Punjabi

Chemistry
Geography
Nafsiyat
Economics

Akhlaqiat
Statistics
Ilm-ul-Taleem
English Grammar & Composition

Pakistan Studies
Computer Science
Mathematics

MATHEMATICS

MULTIPLE CHOICE QUESTIONS WITH ANSWERS

11

MUHAMMAD SALMAN
SHERAZI

M.Sc(Math) , B.ed



	Contents	
UNIT #	Title	Page #
1	Number System	3
2	Sets, Functions and Groups	4
3	Matrices and Determinants	6
4	Quadratic Equations	8
5	Partial Fractions	9
6	Sequence and Series	10
7	Permutation, Combinations and Probability	11
8	Mathematical Induction and Binomial Theorem	13
9	Fundamentals of Trigonometry	13
10	Trigonometric Identities	15
11	Trigonometric Functions and their Graphs	17
12	Application of Trigonometry	18
13	Inverse Trigonometric Functions	19
14	Solution of Trigonometric Equations	21

UNIT # 01 Number Systems

Each question has four possible answer. Tick the correct answer.

- For any complex number z , it is always true that $|z|$ is equal to:
 - $|\bar{z}|$
 - $|-z|$
 - $|- \bar{z}|$
 - ☒ all of these
- The numbers which can be written in the form of $\frac{p}{q}$, $p, q \in \mathbb{Z}$, $q \neq 0$ are :
 - ☒ Rational number
 - Irrational number
 - Complex number
 - Whole number
- A decimal which has a finite numbers of digits in its decimal part is called _____ decimal.
 - ☒ Terminating
 - Non-Terminating
 - Recurring
 - Non recurring
- 5.333 Is
 - Rational
 - Irrational
 - an integer
 - a prime number
- π is
 - Rational
 - ☒ Irrational
 - Natural number
 - None
- $\frac{22}{7}$ is
 - ☒ Rational
 - Irrational
 - an integer
 - a whole number
- Multiplicative inverse of '0' is
 - 0
 - any real number
 - ☒ not defined
 - 1
- If a is any non-zero real number, then multiplicative inverse is
 - $-a$
 - ☒ $\frac{1}{a}$
 - $-\frac{1}{a}$
 - not defined
- For all $a \in \mathbb{R}$, $a = a$ is Property.
 - ☒ Reflexive
 - Symmetric
 - Transitive
 - Trichotomy
- For all $a, b \in \mathbb{R}$, $a = b \Rightarrow b = a$ is called..... property.
 - Reflexive
 - ☒ Symmetric
 - Transitive
 - Trichotomy
- Golden rule of fraction is that for $k \neq 0$, $\frac{a}{b} =$
 - ☒ $\frac{ka}{kb}$
 - $\frac{ab}{l}$
 - $\frac{ka}{b}$
 - $\frac{kb}{b}$
- The set $\{1, -1\}$ possesses closure property w.r. to
 - '+'
 - ☒ '×'
 - $\div$
 - '-'
- If $a < b$ then
 - $a < b$
 - $\frac{1}{a} < \frac{1}{b}$
 - ☒ $\frac{1}{a} > \frac{1}{b}$
 - $a - b > 0$
- The additive identity in set of complex number is
 - ☒ (0,0)
 - (0,1)
 - (1,0)
 - (1,1)
- The multiplicative identity of complex number is
 - (0,0)
 - (0,1)
 - ☒ (1,0)
 - (1,1)
- The modulus of $\bar{z} = a + ib$ is
 - $\sqrt{a+b}$
 - ☒ $\sqrt{a^2 + b^2}$
 - $a - b$
 - $\sqrt{a^2 - b^2}$
- i^{13} equals:
 - ☒ i
 - $-i$
 - 1
 - 1
- The multiplicative inverse of $(4, -7)$ is:
 - $(-\frac{4}{65}, -\frac{7}{65})$
 - $(-\frac{4}{65}, \frac{7}{65})$
 - $(\frac{4}{65}, -\frac{7}{65})$
 - ☒ $(\frac{4}{65}, \frac{7}{65})$
- $(0, 3)(0, 5) =$
 - 15
 - ☒ -15
 - $-8i$
 - $8i$
- $(-1)^{-\frac{21}{2}} =$
 - i
 - ☒ $-i$
 - 1
 - 1
- $\sqrt{3}$ is _____
 - Rational
 - ☒ Irrational
 - Integer
 - Prime
- Product $\sqrt{-2} \times \sqrt{-2}$ is equal to _____
 - 2
 - ☒ 2
 - 0
 - 4
- The imaginary part of the complex number (b, a) is _____
 - ☒ a
 - b
 - ia
 - None of these
- If $z = -1 - i$ then $\bar{z} =$ _____
 - $(-1, -1)$
 - ☒ $(-1, 1)$
 - $(1, -1)$
 - $(1, 1)$
- The property $7.8 + (-7.8) = 0$ is _____
 - Commutative
 - '.' inverse
 - ☒ '+' inverse
 - Associative

26. If $x = 0$ then multiplicative inverse of x is

- (a) 0 (b) 1 (c) $\frac{1}{x}$ (d) ✓ None of these

27. $(-i)^{15} =$ _____

- (a) 1 (b) -1 (c) ✓ i (d) $-i$

28. If z_1 and z_2 are complex numbers then $|z_1 + z_2|$ is _____

- (a) $< |z_1 + z_2|$ (b) ✓ $\leq |z_1| + |z_2|$ (c) $\geq |z_1 + z_2|$ (d) None of these

29. A recurring decimal represents _____

- (a) Real number (b) Natural number (c) ✓ Rational number (d) None of these

30. Every recurring decimal is

- (a) a rational number (b) an irrational number (c) a prime number (d) a whole number

31. Imaginary part of $(-2 + 3i^3)$ is

- (a) -2 (b) 9 (c) 26 (d) ✓ -3

32. The product of two conjugate complex numbers is

- (a) ✓ A real number (b) an imaginary number (c) may be an irrational number (d) not defined

33. $(0, 1)^3$ is equal to:

- (a) ✓ 1 (b) -1 (c) i (d) $-i$

34. If n is an even integer, then $(i)^n$ is equal to:

- (a) ✓ i (b) $-i$ (c) ± 1 (d) 1

35. Factors of $3(x^2 + y^2)$ are:

- (a) $3(x + y)(x - y)$ (b) ✓ $3(x + iy)(x - iy)$ (c) $\sqrt{3}(x + iy)(x - iy)$ (d) None of these

36. Real part of $\frac{2+i}{i}$ is:

- (a) 1 (b) ✓ -2 (c) -1 (d) $\frac{1}{2}$

UNIT # 02 Sets, Functions and Groups

Each question has four possible answer. Tick the correct answer.

1. If $x \in L \cup M$ then

- (a) $x \notin L$ or $x \notin M$ (b) $x \notin L$ or $x \in M$ (c) $x \in L$ or $x \notin M$ (d) ✓ $x \in L$ or $x \in M$

2. Total number of subsets that can be formed from the set $\{x, y, z\}$ is

- (a) 1 (b) ✓ 8 (c) 5 (d) 2

3. If $x \in B' = U - B$ then

- (a) $x \in B$ and $x \in U$ (b) ✓ $x \notin B$ and $x \in U$ (c) $x \notin B$ and $x \notin U$ (d) $x \in B$ and $x \notin U$

4. $L \cup M = L \cap M$ then L is equal to

- (a) ✓ M (b) L (c) $\varnothing$ (d) M'

5. A set is a collection of objects which are

- (a) Well defined (b) ✓ Well defined and distinct (c) identical (d) not defined

6. The set of odd numbers between 1 and 9 are

- (a) $\{1, 3, 5, 7\}$ (b) $\{3, 5, 7, 9\}$ (c) $\{1, 3, 5, 7, 9\}$ (d) ✓ $\{3, 5, 7\}$

7. Every recurring non terminating decimal represents

- (a) ✓ Q (b) Q' (c) R (d) None

8. A diagram which represents a set is called _____

- (a) ✓ Venn's (b) Argand (c) Plane (d) None

9. $R - \{0\}$ is a group w. r. t the binary operation

- (a) $+$ (b) ✓ $\times$ (c) $\div$ (d) $-$

10. In a proposition if $p \rightarrow q$ then $q \rightarrow p$ is called

- (a) Inverse of $p \rightarrow q$ (b) ✓ converse of $p \rightarrow q$ (c) contrapositive of $p \rightarrow q$ (d) None

11. If A and B are disjoint sets, then $A \cap B =$

- (a) 0 (b) 1 (c) 2 (d) ✓ $\varnothing$

12. $p \rightarrow q$ is called converse of

- (a) $\sim pq$ (b) ✓ $p \rightarrow q$ (c) $q \rightarrow p$ (d) $\sim q \rightarrow p$

13. If $A \subseteq B$ then $n(A \cap B) =$

- (a) ✓ $n(A)$ (b) $n(B)$ (c) $n(A) + n(B)$ (d) $n(A) \cdot n(B)$

14. Inverse of any element of a group is:

- (a) Not unique (b) ✓ unique (c) has many inverses (d) none of these

15. Every function is:

- (a) ✓ Relation (b) inverse function (c) one to one (d) none of these

16. The number of all subsets of a set having three elements is:

- (a) 4 (b) 6 (c) ✓ 8 (d) 10

17. The graph of linear function is :

- (a) Circle (b) ✓ straight line (c) parabola (d) triangle

18. Identity element in $(C, .)$ is:

- (a) (0,0) (b) (0,1) (c) ✓ (1,0) (d) (1,1)

19. Squaring a number is a:

- (a) ✓ Unary operation (b) binary operation (c) relation (d) Function

20. Which of the following is true:

- (a) $N \subset Z$ (b) $Z \subset Q$ (c) $Q \subset R$ (d) ✓ all of these

21. Set of integers is a group w.r. t

- (a) ✓ Addition (b) multiplication (c) subtraction (d) division

22. If $A = \varnothing$ then $P(A) =$

- (a) Empty set (b) $\{0\}$ (c) ✓ $\{\varnothing\}$ (d) none of these

23. If A and B are disjoint sets then :

- (a) ✓ $A \cap B = \varnothing$ (b) $A \cap B \neq \varnothing$ (c) $A \subset B$ (d) $A - B = \varnothing$

24. The set of non-zero real numbers w.r. t multiplication is:

- (a) Groupied (b) Semi-group (c) Monoied (d) ✓ Group

25. Which of the following is not a binary operation :

- (a) + (b) $\times$ (c) ✓ $\sqrt{\quad}$ (d) -

26. If $y = \sqrt{x}, x \geq 0$ is a function, then its inverse is:

- (a) A line (b) a parabola (c) a point (d) ✓ not a function

27. An onto function is also called:

- (a) Injective (b) ✓ Surjective (c) Bijective (d) Inverse

28. A $(1 - 1)$ function is also called:

- (a) ✓ Injective (b) Surjective (c) Bijective (d) Inverse

29. For the propositions p and q , $(p \wedge q) \rightarrow p$ is:

- (a) ✓ Tautology (b) Absurdity (c) contingency (d) None of these

30. For the propositions p and q , $p \rightarrow (p \vee q)$ is:

- (a) ✓ Tautology (b) Absurdity (c) Contingency (d) None of these

31. If set A has 2 elements and B has 4 elements, then number of elements in $A \times B$ is :

- (a) 6 (b) ✓ 8 (c) 16 (d) None of these

32. Inverse of a line is :

- (a) ✓ A line (b) a parabola (c) a point (d) not defined

33. The function $f = \{(x, y), y = x\}$ is :

- (a) ✓ Identity function (b) Null function (c) not a function (d) similar function

34. If $y = \sqrt{x}, x \geq 0$ is a function, then its inverse is :

- (a) A line (b) a parabola (c) a point (d) ✓ not defined

35. Truth set of a tautology is

- (a) ✓ Universal set (b) $\varnothing$ (c) True (d) False

36. A compound proposition which is always true is called:

- (a) ✓ Tautology (b) contradiction (c) absurdity (d) contingency

37. A compound proposition which is always neither true nor false is called:

- (a) Tautology (b) contradiction (c) absurdity (d) ✓ contingency

38. A compound proposition which is always wrong is called:

- (a) Tautology (b) ✓ contradiction (c) absurdity (d) contingency

39. If $S = \{ \}$, then order of set S is:

- (a) ✓ 0 (b) 1 (c) Infinite set (d) not defined

40. The symbol which is used to denote negation of a proposition is

- (a) ✓ $\sim$ (b) $\rightarrow$ (c) $\wedge$ (d) $\vee$

41. Which of the following is true:

- (a) $\varnothing - A = \varnothing$ (b) $A \cup A = A$ (c) $A \cap A = A$ (d) ✓ All of these

42. Which of the following is true:

- (a) $A \cup \varnothing = A$ (b) $A \cap \varnothing = \varnothing$ (c) $A - \varnothing = A$ (d) ✓ All of these

43. If $B \subseteq A$, then $n(B - A)$ is equal to:

- (a) $n(A)$ (b) $n(B)$ (c) $n(A \cap B)$ (d) ✓ 0

UNIT # 03 Matrices and Determinants

Each question has four possible answer. Tick the correct answer.

- A rectangular array of numbers enclosed by a square brackets is called:
 - ✓ Matrix
 - Row
 - Column
 - Determinant
- The horizontal lines of numbers in a matrix are called:
 - Columns
 - ✓ Rows
 - Column matrix
 - Row matrix
- The vertical lines of numbers in a matrix are called:
 - ✓ Columns
 - Rows
 - Column matrix
 - Row matrix
- If a matrix A has m rows and n columns, then order of A is :
 - ✓ $m \times n$
 - $n \times m$
 - $m + n$
 - m^n
- The element a_{ij} of any matrix A is present in:
 - ✓ i^{th} row and j^{th} column
 - i^{th} column and j^{th} row
 - $(i + j)^{th}$ row and column
 - $(i + j)^{th}$ row and column
- A matrix A is called real if all a_{ij} are
 - ✓ Real numbers
 - complex numbers
 - 0
 - 1
- If a matrix A has only one row then it is called:
 - ✓ Row vector
 - Column vector
 - square matrix
 - Rectangle matrix
- If a matrix A has only one column then it is called:
 - Row vector
 - ✓ Column vector
 - square matrix
 - Rectangle matrix
- If a matrix A has same number of rows and columns then A is called:
 - Row vector
 - Column vector
 - ✓ square matrix
 - Rectangular matrix
- If a matrix A has different number of rows and columns then A is called:
 - Row vector
 - Column vector
 - square matrix
 - ✓ Rectangular matrix
- For the square matrix $A = [a_{ij}]_{n \times n}$ then $a_{11}, a_{22}, a_{33} \dots a_{nn}$ are:
 - ✓ Main diagonal
 - primary diagonal
 - proceeding diagonal
 - secondary diagonal
- For a square matrix $A = [a_{ij}]$ if all $a_{ij} = 0, i \neq j$ and all $a_{ij} = k$ (non-zero) for $i = j$ then A is called:
 - Diagonal matrix
 - ✓ Scalar Matrix
 - Scalar Matrix
 - Null matrix
- For a square matrix $A = [a_{ij}]$ if all $a_{ij} = 0, i \neq j$ and at least $a_{ij} \neq 0, i = j$ then A is called:
 - ✓ Diagonal matrix
 - Scalar Matrix
 - Scalar Matrix
 - Null matrix
- The matrix $[7]$ is :
 - Square matrix
 - row matrix
 - column matrix
 - ✓ all of these
- If a matrix of order $m \times n$ then the matrix of order $n \times m$ is :
 - ✓ Transpose of A
 - Inverse of A
 - Main diagonal of A
 - Adjoint of A
- For any two matrices A and $B, (A + B)^t =$
 - $B^t A^t$
 - $A^t B^t$
 - ✓ $A^t + B^t$
 - $B^t + A^t$
- Let A be a matrix and n is an integer then $A + A + A + \dots$ to n terms
 - ✓ nA
 - $n + A$
 - $n - A$
 - A^n
- If order of A is $m \times n$ and order of B is $n \times q$ then order of AB is
 - $m \times n$
 - $n \times m$
 - $n \times q$
 - ✓ $m \times q$
- If $adj A = \begin{bmatrix} -1 & -2 \\ 3 & 4 \end{bmatrix}$ then matrix A is
 - $\begin{bmatrix} -1 & -2 \\ 4 & 3 \end{bmatrix}$
 - $\begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix}$
 - $\begin{bmatrix} -4 & 3 \\ -2 & -1 \end{bmatrix}$
 - ✓ $\begin{bmatrix} 4 & 2 \\ -3 & -1 \end{bmatrix}$
- If A is non-singular matrix then $A^{-1} =$
 - ✓ $\frac{1}{|A|} adj A$
 - $-\frac{1}{|A|} adj A$
 - $\frac{|A|}{adj A}$
 - $\frac{1}{|A| adj A}$
- If $AX = B$ then X is equal to:
 - AB
 - ✓ $A^{-1}B$
 - $B^{-1}A$
 - BA
- Inverse of a matrix exists if it is:
 - Singular
 - Null
 - Rectangular
 - ✓ Non-singular
- Which of the property does not hold matrix multiplication?
 - Associative
 - ✓ Commutative
 - Closure
 - Inverse
- For any matrix A , it is always true that
 - $A = A^t$
 - $-A = \bar{A}$
 - ✓ $|A| = |A^t|$
 - $A^{-1} = \frac{1}{A}$
- If all entries of a square matrix of order 3 is multiplied by k , then value of $|kA|$ is equal to:
 - $k|A|$
 - $k^2|A|$
 - ✓ $k^3|A|$
 - $|A|$

26. For a non-singular matrix it is true that :

- (a) $(A^{-1})^{-1} = A$ (b) $(A^t)^t = A$ (c) $\bar{\bar{A}} = A$ (d) ✓ all of these

27. For any non-singular matrices A and B it is true that:

- (a) $(AB)^{-1} = B^{-1}A^{-1}$ (b) $(AB)^t = B^tA^t$ (c) $AB \neq BA$ (d) ✓ all of these

28. A square matrix $A = [a_{ij}]$ for which $a_{ij} = 0, i > j$ then A is called:

- (a) ✓ Upper triangular (b) Lower triangular (c) Symmetric (d) Hermitian

29. A square matrix $A = [a_{ij}]$ for which $a_{ij} = 0, i < j$ then A is called:

- (a) Upper triangular (b) ✓ Lower triangular (c) Symmetric (d) Hermitian

30. Any matrix A is called singular if:

- (a) ✓ $|A| = 0$ (b) $|A| \neq 0$ (c) $A^t = A$ (d) $AA^{-1} = I$

31. Which of the following Sets is a field.

- (a) R (b) Q (c) C (d) ✓ all of these

32. Which of the following Sets is not a field.

- (a) R (b) Q (c) C (d) ✓ Z

33. A square matrix A is symmetric if:

- (a) ✓ $A^t = A$ (b) $A^t = -A$ (c) $(\bar{A})^t = A$ (d) $(\bar{A})^t = -A$

34. A square matrix A is skew symmetric if:

- (a) $A^t = A$ (b) ✓ $A^t = -A$ (c) $(\bar{A})^t = A$ (d) $(\bar{A})^t = -A$

35. A square matrix A is Hermitian if:

- (a) $A^t = A$ (b) $A^t = -A$ (c) ✓ $(\bar{A})^t = A$ (d) $(\bar{A})^t = -A$

36. A square matrix A is skew- Hermitian if:

- (a) $A^t = A$ (b) $A^t = -A$ (c) $(\bar{A})^t = A$ (d) ✓ $(\bar{A})^t = -A$

37. The main diagonal elements of a skew symmetric matrix must be:

- (a) 1 (b) ✓ 0 (c) any non-zero number (d) any complex number

38. The main diagonal elements of a skew hermitian matrix must be:

- (a) 1 (b) ✓ 0 (c) any non-zero number (d) any complex number

39. In echelon form of matrix, the first non zero entry is called:

- (a) ✓ Leading entry (b) first entry (c) preceding entry (d) Diagonal entry

40. The additive inverse of a matrix exist only if it is:

- (a) Singular (b) non singular (c) null matrix (d) ✓ any matrix of order $m \times n$

41. The multiplicative inverse of a matrix exist only if it is:

- (a) Singular (b) ✓ non singular (c) null matrix (d) any matrix of order $m \times n$

42. The number of non zero rows in echelon form of a matrix is called:

- (a) Order of matrix (b) Rank of matrix (c) leading (d) leading row

43. If A is any square matrix then $A + A^t$ is a

- (a) ✓ Symmetric (b) skew symmetric (c) hermitian (d) skew hermitian

44. If A is any square matrix then $A - A^t$ is a

- (a) Symmetric (b) ✓ skew symmetric (c) hermitian (d) skew hermitian

45. If A is any square matrix then $A + (\bar{A})^t$ is a

- (a) Symmetric (b) skew symmetric (c) ✓ hermitian (d) skew hermitian

46. If A is any square matrix then $A + (\bar{A})^t$ is a

- (a) Symmetric (b) skew symmetric (c) hermitian (d) ✓ skew hermitian

47. If A is symmetric (Skew symmetric), then A^2 must be

- (a) Singular (b) non singular (c) ✓ symmetric (d) non trivial solution

48. In a homogeneous system of linear equations , the solution (0,0,0) is:

- (a) ✓ Trivial solution (b) non trivial solution (c) exact solution (d) anti symmetric

49. If $AX = O$ then $X =$

- (a) I (b) ✓ O (c) A^{-1} (d) Not possible

50. If the system of linear equations have no solution at all, then it is called a/an

- (a) Consistent system (b) ✓ Inconsistent system (c) Trivial System (d) Non Trivial System

51. The value of λ for which the system $x + 2y = 4; 2x + \lambda y = -3$ does not possess the unique solution

- (a) ✓ 4 (b) -4 (c) ± 4 (d) any real number

52. If the system $x + 2y = 0; 2x + \lambda y = 0$ has non-trivial solution, then λ is:

- (a) ✓ 4 (b) -4 (c) ± 4 (d) any real number

53. The inverse of unit matrix is:

- (a) ✓ Unit (b) Singular (c) Skew Symmetric (d) rectangular

54. Transpose of a row matrix is:

- (a) Diagonal matrix (b) zero matrix (c) ✓ column matrix (d) scalar matrix

55. If $\begin{vmatrix} x & 4 \\ 5 & 10 \end{vmatrix} = 0 \Rightarrow x$ equals

- (a) ✓ 2 (b) 4 (c) 6 (d) 8

UNIT # 04 Quadratic Equations

Each question has four possible answer. Tick the correct answer.

1. The equation $ax^2 + bx + c = 0$ will be quadratic if:

- (a) $a = 0, b \neq 0$ (b) ✓ $a \neq 0$ (c) $a = b = 0$ (d) $b =$ any real number

2. Solution set of the equation $x^2 - 4x + 4 = 0$ is:

- (a) $\{2, -2\}$ (b) ✓ $\{2\}$ (c) $\{-2\}$ (d) $\{4, -4\}$

3. The quadratic formula for solving the equation $ax^2 + bx + c = 0; a \neq 0$ is

- (a) ✓ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ (b) $x = \frac{-b \pm \sqrt{a^2 - 4ac}}{2b}$ (c) $x = \frac{-a \pm \sqrt{b^2 - 4ac}}{2}$ (d) None of these

4. To convert $ax^{2n} + bx^n + c = 0 (a \neq 0)$ into quadratic form, the correct substitution is:

- (a) ✓ $y = x^n$ (b) $x = y^n$ (c) $y = x^{-n}$ (d) $y = \frac{1}{x}$

5. The equation in which variable occurs in exponent, called:

- (a) Exponential function (b) Quadratic equation (c) Reciprocal equation (d) ✓ Exponential equation

6. To convert $4^{1+x} + 4^{1-x} = 10$ into quadratic, the substitution is:

- (a) $y = x^{1-x}$ (b) $y = 4^{1+x}$ (c) $y = 4^x$ (d) $y = 4^{-x}$

7. The equations involving redical expressions of the variable are called:

- (a) Reciprocal equations (b) ✓ Redical equations (c) Quadratic functions (d) exponential equations

8. The cube roots of unity are :

- (a) ✓ $1, \frac{-1+\sqrt{3}i}{2}, \frac{-1-\sqrt{3}i}{2}$ (b) $1, \frac{1+\sqrt{3}i}{2}, \frac{1+\sqrt{3}i}{2}$ (c) $-1, \frac{-1+\sqrt{3}i}{2}, \frac{-1+\sqrt{3}i}{2}$ (d) $-1, \frac{1+\sqrt{3}i}{2}, \frac{1+\sqrt{3}i}{2}$

9. Sum of all cube roots of 64 is :

- (a) ✓ 0 (b) 1 (c) 64 (d) -64

10. Product of cube roots of -1 is:

- (a) 0 (b) -1 (c) ✓ 1 (d) None

11. $16\omega^8 + 16\omega^4 =$

- (a) 0 (b) ✓ -16 (c) 16 (d) -1

12. The sum of all four fourth roots of unity is:

- (a) Unity (b) ✓ 0 (c) -1 (d) None

13. The product of all four fourth roots of unity is:

- (a) Unity (b) 0 (c) ✓ -1 (d) None

14. The sum of all four fourth roots of 16 is:

- (a) 16 (b) -16 (c) ✓ 0 (d) 1

15. The complex cube roots of unity are..... each other.

- (a) Additive inverse (b) Equal to (c) ✓ Conjugate (d) None of these

16. The complex cube roots of unity are..... each other.

- (a) Multiplicative inverse (b) reciprocal (c) square (d) ✓ None of these

17. The complex fourth roots of unity are of each other.

- (a) ✓ Additive inverse (b) equal to (c) square of (d) None of these

18. If sum of all cube roots of unity is equal to $x^2 + 1$, then x is equal to:

- (a) -1 (b) 0 (c) ✓ $\pm i$ (d) 1

19. If product of all cube roots of unity is equal to $p^2 + 1$, then p is equal to:

- (a) -1 (b) ✓ 0 (c) $\pm i$ (d) 1

20. The expression $x^2 + \frac{1}{x} - 3$ is polynomial of degree:

- (a) 2 (b) 3 (c) 1 (d) ✓ not a polynomial

21. If $f(x)$ is divided by $x - a$, then dividend = (Divisor)(.....)+ Remainder.

- (a) Divisor (b) Dividend (c) ✓ Quotient (d) $f(a)$

22. If $f(x)$ is divided by $x - a$ by remainder theorem then remainder is:

- (a) ✓ $f(a)$ (b) $f(-a)$ (c) $f(a) + R$ (d) $x - a = R$

23. The polynomial $(x - a)$ is a factor of $f(x)$ if and only if

- (a) ✓ $f(a) = 0$ (b) $f(a) = R$ (c) Quotient = R (d) $x = -a$

24. $x - 2$ is a factor of $x^2 - kx + 4$, if k is:

- (a) 2 (b) ✓ 4 (c) 8 (d) -4

25. If $x = -2$ is the root of $kx^4 - 13x^2 + 36 = 0$, then $k =$

- (a) 2 (b) -2 (c) 1 (d) ✓ -1
26. $x + a$ is a factor of $x^n + a^n$ when n is
 (a) Any integer (b) any positive integer (c) ✓ any odd integer (d) any real number
27. $x - a$ is a factor of $x^n - a^n$ when n is
 (a) ✓ Any integer (b) any positive integer (c) any odd integer (d) any real number
28. Sum of roots of $ax^2 - bx - c = 0$ is ($a \neq 0$)
 (a) ✓ $\frac{b}{a}$ (b) $-\frac{b}{a}$ (c) $\frac{c}{a}$ (d) $-\frac{c}{a}$
29. Sum of roots of $ax^2 - bx - c = 0$ is ($a \neq 0$)
 (a) $\frac{b}{a}$ (b) $-\frac{b}{a}$ (c) $\frac{c}{a}$ (d) ✓ $-\frac{c}{a}$
30. If 2 and -5 are roots of a quadratic equation, then equation is:
 (a) $x^2 - 3x - 10 = 0$ (b) $x^2 - 3x + 10 = 0$ (c) ✓ $x^2 + 3x - 10 = 0$ (d) $x^2 + 3x + 10 = 0$
31. If α and β are the roots of $3x^2 - 2x + 4 = 0$, then the value of $\alpha + \beta$ is:
 (a) ✓ $\frac{2}{3}$ (b) $-\frac{2}{3}$ (c) $\frac{4}{3}$ (d) $-\frac{4}{3}$
32. If roots of $ax^2 + bx + c = 0$, ($a \neq 0$) are real, then
 (a) $\text{Disc} \geq 0$ (b) $\text{Disc} < 0$ (c) $\text{Disc} \neq 0$ (d) $\text{Disc} \leq 0$
33. If roots of $ax^2 + bx + c = 0$, ($a \neq 0$) are complex, then
 (a) ✓ $\text{Disc} \geq 0$ (b) ✓ $\text{Disc} < 0$ (c) $\text{Disc} \neq 0$ (d) $\text{Disc} \leq 0$
34. If roots of $ax^2 + bx + c = 0$, ($a \neq 0$) are equal, then
 (a) ✓ $\text{Disc} = 0$ (b) $\text{Disc} < 0$ (c) $\text{Disc} \neq 0$ (d) None of these
35. Graph of quadratic equation is:
 (a) Straight line (b) Circle (c) square (d) ✓ Parabola
36. $\omega^{28} + \omega^{29} + 1 =$
 (a) ✓ 0 (b) 1 (c) -1 (d) ω
37. Synthetic division is a process of:
 (a) Addition (b) multiplication (c) subtraction (d) ✓ division
38. Degree of quadratic equation is:
 (a) 0 (b) 1 (c) ✓ 2 (d) 3
39. Basic techniques for solving quadratic equations are
 (a) 1 (b) 2 (c) ✓ 3 (d) 4
40. $16\omega^4 + 16\omega^8 =$
 (a) 0 (b) ✓ -16 (c) 16 (d) -1

UNIT # 05 Partial Fractions



Each question has four possible answer. Tick the correct answer.

1. An open sentence formed by using sign of "=" is called a/an
 (a) ✓ Equation (b) Formula (c) Rational fraction (d) Theorem
2. If an equation is true for all values of the variable, then it is called:
 (a) a conditional equation (b) ✓ an identity (c) proper rational fraction (d) All of these
3. $(x + 3)(x + 4) = x^2 + 7x + 12$ is a/an:
 (a) Conditional equation (b) ✓ an identity (c) proper rational fraction (d) a formula
4. The quotient of two polynomials $\frac{P(x)}{Q(x)}$, $Q(x) \neq 0$ is called :
 (a) ✓ Rational fraction (b) Irrational fraction (c) Partial fraction (d) Proper fraction
5. A fraction $\frac{P(x)}{Q(x)}$, $Q(x) \neq 0$ is called proper fraction if :
 (a) ✓ Degree of $P(x) < \text{Degree of } Q(x)$ (b) Degree of $P(x) = \text{Degree of } Q(x)$
 (c) Degree of $P(x) > \text{Degree of } Q(x)$ (d) Degree of $P(x) \geq \text{Degree of } Q(x)$
6. A fraction $\frac{P(x)}{Q(x)}$, $Q(x) \neq 0$ is called proper fraction if :
 (a) Degree of $P(x) < \text{Degree of } Q(x)$ (b) Degree of $P(x) = \text{Degree of } Q(x)$
 (c) Degree of $P(x) > \text{Degree of } Q(x)$ (d) ✓ Degree of $P(x) \geq \text{Degree of } Q(x)$
7. A mixed form of fraction is :
 (a) An integer+ improper fraction (b) a polynomial+improper fraction
 (c) ✓ a polynomial+proper fraction (d) a polynomial+rational fraction
8. When a rational fraction is separated into partial fractions, then result is always :
 (a) A conditional equations (b) ✓ an identity (c) a partial fraction (d) an improper fraction

9. The number of Partial fraction of $\frac{x^3}{x(x+1)(x^2-1)}$ are:
 (a) 2 (b) 3 (c) ✓ 4 (d) None of these
10. The number of Partial fraction of $\frac{x^5}{x(x+1)(x^2-4)}$ are:
 (a) 2 (b) 3 (c) 4 (d) ✓ 6
11. Partial fractions of $\frac{1}{x^2-1}$ are:
 (a) $\frac{1}{2(x-1)} + \frac{1}{2(x+1)}$ (b) ✓ $\frac{1}{2(x-1)} - \frac{1}{2(x+1)}$ (c) $-\frac{1}{2(x-1)} + \frac{1}{2(x+1)}$ (d) $-\frac{1}{2(x-1)} - \frac{1}{2(x+1)}$
12. Conditional equation $2x + 3 = 0$ holds when x is equal to:
 (a) ✓ $-\frac{3}{2}$ (b) $\frac{3}{2}$ (c) $\frac{1}{3}$ (d) 1
13. Which is a reducible factor:
 (a) $x^3 - 6x^2 + 8x$ (b) $x^2 + 16x$ (c) $x^2 + 5x - 6$ (d) ✓ all of these
14. A quadratic factor which cannot written as a product of linear factors with real coefficients is called:
 (a) ✓ An irreducible factor (b) reducible factor (c) an irrational factor (d) an improper factor
15. $\frac{9x^2}{x^3-1}$ is an
 (a) Improper fraction (b) ✓ Proper fraction (c) Polynomial (d) equation

UNIT # 06 Sequence and Series

Each question has four possible answer. Tick the correct answer.

1. An arrangement of numbers according to some definite rule is called:
 (a) ✓ Sequence (b) Combination (c) Series (d) Permutation
2. A sequence is also known as:
 (a) Real sequence (b) ✓ Progression (c) Arrangement (d) Complex sequence
3. A sequence is function whose domain is
 (a) Z (b) N (c) Q (d) ✓ R
4. As sequence whose range is R i. e., set of real numbers is called:
 (a) ✓ Real sequence (b) Imaginary sequence (c) Natural sequence (d) Complex sequence
5. If $a_n = \{n + (-1)^n\}$, then $a_{10} =$
 (a) 10 (b) ✓ 11 (c) 12 (d) 13
6. The last term of an infinite sequence is called :
 (a) n th term (b) a_n (c) last term (d) ✓ does not exist
7. The next term of the sequence 1, 2, 12, 40, ... is
 (a) ✓ 112 (b) 120 (c) 124 (d) None of these
8. A sequence $\{a_n\}$ in which $a_n - a_{n-1}$ is the same number for all $n \in N, n > 1$ is called:
 (a) ✓ A.P (b) G.P (c) H.P (d) None of these
9. n th term of an A.P is $3n - 1$ then 10th term is :
 (a) 9 (b) ✓ 29 (c) 12 (d) cannot determined
10. n th term of the series $\left(\frac{1}{3}\right)^2 + \left(\frac{5}{2}\right)^2 + \left(\frac{7}{3}\right)^2 + \dots$
 (a) ✓ $\left(\frac{2n-1}{3}\right)^2$ (b) $\left(\frac{2n-1}{3}\right)^2$ (c) $\left(\frac{2n}{3}\right)^2$ (d) cannot determined
11. If a_{n-1}, a_n, a_{n+1} are in A.P, then a_n is
 (a) ✓ A.M (b) G.M (c) H.M (d) Mid point
12. Arithmetic mean between c and d is:
 (a) ✓ $\frac{c+d}{2}$ (b) $\frac{c+d}{2cd}$ (c) $\frac{2cd}{c+d}$ (d) $\frac{2}{c+d}$
13. The arithmetic mean between $\sqrt{2}$ and $3\sqrt{2}$ is:
 (a) $4\sqrt{2}$ (b) ✓ $\frac{4}{\sqrt{2}}$ (c) $\sqrt{2}$ (d) none of these
14. The sum of terms of a sequence is called:
 (a) Partial sum (b) ✓ Series (c) Finite sum (d) none of these
15. Forth partial sum of the sequence $\{n^2\}$ is called:
 (a) 16 (b) ✓ $1+4+9+16$ (c) 8 (d) $1+2+3+4$
16. Sum of n -term of an Arithmetic series S_n is equal to:
 (a) ✓ $\frac{n}{2}[2a + (n-1)d]$ (b) $\frac{n}{2}[a + (n-1)d]$ (c) $\frac{n}{2}[2a + (n+1)d]$ (d) $\frac{n}{2}[2a + l]$
17. For any G. P., the common ratio r is equal to:
 (a) $\frac{a_n}{a_{n+1}}$ (b) $\frac{a_{n-1}}{a_n}$ (c) ✓ $\frac{a_n}{a_{n-1}}$ (d) $a_{n+1} - a_n, n \in N, n > 1$

18. No term of a G.P., is:

- (a) ☒ 0 (b) 1 (c) negative (d) imaginary number

19. The general term of a G.P., is :

- (a) ☒ $a_n = ar^{n-1}$ (b) $a_n = ar^n$ (c) $a_n = ar^{n+1}$ (d) None of these

20. The sum of infinite geometric series is valid if

- (a) $|r| > 1$ (b) $|r| = 1$ (c) $|r| \geq 1$ (d) ☒ $|r| < 1$

21. For the series $1 + 5 + 25 + 125 + \dots + \infty$, the sum is

- (a) -4 (b) 4 (c) $\frac{1-5^n}{-4}$ (d) ☒ not defined

22. An infinite geometric series is convergent if

- (a) $|r| > 1$ (b) $|r| = 1$ (c) $|r| \geq 1$ (d) ☒ $|r| < 1$

23. An infinite geometric series is divergent if

- (a) $|r| < 1$ (b) $|r| \neq 1$ (c) $r = 0$ (d) ☒ $|r| > 1$

24. If sum of series is defined then it is called:

- (a) ☒ Convergent series (b) Divergent series (c) finite series (d) Geometric series

25. If sum of series is not defined then it is called:

- (a) Convergent series (b) ☒ Divergent series (c) finite series (d) Geometric series

26. The interval in which series $1 + 2x + 4x^2 + 8x^3 + \dots$ is convergent if :

- (a) $-2 < x < 2$ (b) ☒ $-\frac{1}{2} < x < \frac{1}{2}$ (c) $|2x| > 1$ (d) $|x| < 1$

27. If the reciprocal of the terms a sequence form an A.P., then it is called:

- (a) ☒ H.P. (b) G.P. (c) A.P. (d) sequence

28. The n th term of $\frac{1}{2}, \frac{1}{5}, \frac{1}{8}, \dots$ is

- (a) ☒ $\frac{1}{3n-1}$ (b) $3n-1$ (c) $2n+1$ (d) $\frac{1}{3n+1}$

29. Harmonic mean between 2 and 8 is:

- (a) ☒ 5 (b) $\frac{16}{5}$ (c) ± 4 (d) $\frac{5}{16}$

30. If A, G and H are Arithmetic, Geometric and Harmonic means between two positive numbers then

- (a) $G^2 = AH$ (b) ☒ A, G, H are in G.P. (c) $A > G > H$ (d) all of these

31. If A, G and H are Arithmetic, Geometric and Harmonic means between two negative numbers then

- (a) $G^2 = AH$ (b) A, G, H are in G.P. (c) $A < G < H$ (d) ☒ all of these

32. If a and b are two positive number then

- (a) $A < G < H$ (b) ☒ $A > G > H$ (c) $A = G = H$ (d) $A \geq G \geq H$

33. If a and b are two negative number then

- (a) ☒ $A < G < H$ (b) $A > G > H$ (c) $A = G = H$ (d) $A \geq G \geq H$

34. If $\frac{a^{n+1}+b^{n+1}}{a^n+b^n}$ is A.M between a and b then n is equal to:

- (a) ☒ 0 (b) -1 (c) 1 (d) $\frac{1}{2}$

35. If $\frac{a^n+b^n}{a^{n-1}+b^{n-1}}$ is G.M between a and b then n is equal to:

- (a) 0 (b) -1 (c) 1 (d) ☒ $\frac{1}{2}$

36. If $\frac{a^{n+1}+b^{n+1}}{a^n+b^n}$ is H.M between a and b then n is equal to:

- (a) 0 (b) ☒ -1 (c) 1 (d) $\frac{1}{2}$

37. If $S_n = (n+1)^2$, then S_{2n} is equal to:

- (a) $2n+1$ (b) ☒ $4n^2 + 4n + 1$ (c) $(2n-1)^2$ (d) cannot be determined

38. $\sum n^3 =$

- (a) $\frac{n(n+1)}{2}$ (b) $\frac{n(n+1)(n+2)}{6}$ (c) ☒ $\frac{n^2(n+1)^2}{4}$ (d) $\frac{n(n+1)^2}{2}$

UNIT # 07 Permutation, Combination and Probability

Each question has four possible answer. Tick the correct answer.

1. ${}^{20}P_3 =$

- (a) 6890 (b) 6810 (c) ☒ 6840 (d) 6880

2. If ${}^{n}P_2 = 30$ then n =

- (a) 4 (b) 5 (c) ☒ 6 (d) 10

3. The number of diagonals in 10-sided figure is
 (a) 10 (b) 10_{C_2} (c) $\checkmark 10_{C_2} - 10$ (d) 45
4. $n_{C_0} =$
 (a) 0 (b) $\checkmark 1$ (c) n (d) $n!$
5. How many arrangement of the word "MATHEMATICS" can be made
 (a) $11!$ (b) $\binom{11}{3,2,1,1,1,1,1}$ (c) $\checkmark \binom{11}{2,2,2,1,1,1,1}$ (d) None
6. If n is negative then $n!$ is
 (a) 1 (b) 0 (c) unique (d) $\checkmark$ Not defined
7. $n - 1_{C_r} + n - 1_{C_{r-1}}$ equals
 (a) $\checkmark n_{C_r}$ (b) $n_{C_{r-1}}$ (c) $n - 1_{C_r}$ (d) $n + 1_{C_r}$
8. How many signals can be given by 5 flags of different colors, using 3 at a time
 (a) 120 (b) $\checkmark 60$ (c) 24 (d) 15
9. $n_{P_n} =$
 (a) 1 (b) n (c) $\checkmark n!$ (d) None
10. $\frac{8!}{7!} =$
 (a) $\checkmark 8$ (b) 7 (c) 56 (d) $\frac{8}{7}$
11. If an event A can occur in p ways and B can occur q ways, then number of ways that both events occur is:
 (a) $p + q$ (b) $\checkmark p \cdot q$ (c) $(pq)!$ (d) $(p + q)!$
12. If $\binom{n}{12} = \binom{n}{8}$ then the value of n ,
 (a) 15 (b) 16 (c) 18 (d) $\checkmark 20$
13. Probability of non-occurrence of an event E is equal to :
 (a) $\checkmark 1 - P(E)$ (b) $P(E) + \frac{n(S)}{n(E)}$ (c) $\frac{n(S)}{n(E)}$ (d) $1 + P(E)$
14. For independent events $P(A \cap B) =$
 (a) $P(A) + P(B)$ (b) $P(A) - P(B)$ (c) $\checkmark P(A) \cdot P(B)$ (d) $\frac{P(A)}{P(B)}$
15. A card is drawn from a deck of 52 playing cards. The probability of card that it is an ace card is:
 (a) $\frac{2}{13}$ (b) $\frac{4}{13}$ (c) $\checkmark \frac{1}{13}$ (d) $\frac{17}{13}$
16. Four persons wants to sit in a circular sofa, the total ways are:
 (a) 24 (b) $\checkmark 6$ (c) 4 (d) None of these
17. Two teams A and B are playing a match, the probability that team A does not lose is:
 (a) $\frac{1}{3}$ (b) $\frac{2}{3}$ (c) 1 (d) 0
18. Let $S = \{1, 2, 3, \dots, 10\}$ the probability that a number is divided by 4 is :
 (a) $\frac{2}{5}$ (b) $\checkmark \frac{1}{5}$ (c) $\frac{1}{10}$ (d) $\frac{1}{2}$
19. A die is rolled, the probability of getting 3 or 5 is:
 (a) $\frac{2}{3}$ (b) $\checkmark \frac{15}{36}$ (c) $\frac{15}{36}$ (d) $\frac{1}{36}$
20. A coin is tossed 5 times, then $n(S)$ is equal to:
 (a) $\checkmark 32$ (b) 25 (c) 10 (d) 20
21. The number of ways for sitting 4 persons in a train on a straight sofa is:
 (a) $\checkmark 24$ (b) 6 (c) 4 (d) None of these
22. Sample space for tossing a coin is:
 (a) $\{H\}$ (b) $\{T\}$ (c) $\{H, H\}$ (d) $\checkmark \{H, T\}$
23. If $P(E) = \frac{7}{12}$, $n(S) = 8400$, $n(E) =$
 (a) 108 (b) $\checkmark 4900$ (c) 144 (d) 14400
24. In a permutation n_{P_r} or $P(n, r)$, it is always true that
 (a) $\checkmark n \geq r$ (b) $n < r$ (c) $n \leq r$ (d) $n < 0, r < 0$
25. If an event always occurs, then it is called:
 (a) Null Event (b) Possible Event (c) $\checkmark$ Certain event (d) Independent Event
26. If E is a certain event, then
 (a) $P(E) = 0$ (b) $\checkmark P(E) = 1$ (c) $0 < P(E) < 1$ (d) $P(E) > 1$
27. If E is an impossible event, then
 (a) $\checkmark P(E) = 0$ (b) $P(E) = 1$ (c) $P(E) \neq 0$ (d) $0 < P(E) < 1$

28. Non occurrence of an event E is denoted by:

- (a) $\sim E$ (b) $\checkmark \bar{E}$ (c) E^c (d) All of these

UNIT # 08 Mathematical Induction and Binomial Theorem

Each question has four possible answer. Tick the correct answer.

- The statement $4^n + 3^n + 4$ is true when :
 (a) $n = 0$ (b) $n = 1$ (c) $\checkmark n \geq 2$ (d) n is any +iv integer
- The number of terms in the expansion of $(a + b)^n$ are:
 (a) n (b) $\checkmark n + 1$ (c) 2^n (d) 2^{n-1}
- Middle term/s in the expansion of $(a - 3x)^{14}$ is/are :
 (a) T_7 (b) $\checkmark T_8$ (c) $T_6 \& T_7$ (d) $T_7 \& T_8$
- The coefficient of the last term in the expansion of $(2 - x)^7$ is :
 (a) 1 (b) $\checkmark -1$ (c) 7 (d) -7
- $\binom{2n}{0} + \binom{2n}{1} + \binom{2n}{2} + \dots + \binom{2n}{2n}$ is equal to:
 (a) 2^n (b) $\checkmark 2^{2n}$ (c) 2^{2n-1} (d) 2^{2n+1}
- $1 + x + x^2 + x^3 + \dots$
 (a) $(1 + x)^{-1}$ (b) $\checkmark (1 - x)^{-1}$ (c) $(1 + x)^{-2}$ (d) $(1 - x)^{-2}$
- The middle term in the expansion of $(a + b)^n$ is $\binom{n}{2} + 1$; then n is
 (a) Odd (b) $\checkmark$ even (c) prime (d) none of these
- The number of terms in the expansion of $(a + b)^{20}$ is:
 (a) 18 (b) 20 (c) $\checkmark 21$ (d) 19
- The expansion $(1 - 4x)^{-2}$ is valid if:
 (a) $\checkmark |x| < \frac{1}{4}$ (b) $|x| > \frac{1}{4}$ (c) $-1 < x < 1$ (d) $|x| < -1$
- The statement $3^n < n!$ is true, when
 (a) $n = 2$ (b) $n = 4$ (c) $n = 6$ (d) $\checkmark n > 6$
- General term in the expansion of $(a + b)^n$ is:
 (a) $\binom{n+1}{r} a^{n-r} x^r$ (b) $\checkmark \binom{n}{r-1} a^{n-r} x^r$ (c) $\binom{n}{r+1} a^{n-r} x^r$ (d) $\binom{n}{r} a^{n-r} x^r$
- The method of induction was given by Francesco who lived from:
 (a) $\checkmark 1494-1575$ (b) 1500-1575 (c) 1498-1575 (d) 1494-1570

UNIT # 09 Fundamentals of Trigonometry

Each question has four possible answer. Tick the correct answer.

- Two rays with a common starting point form:
 (a) Triangle (b) $\checkmark$ Angle (c) Radian (d) Minute
- The common starting point of two rays is called:
 (a) Origin (b) Initial Point (c) $\checkmark$ Vertex (d) All of these
- If the rotation of the angle is counter clock wise, then angle is:
 (a) Negative (b) $\checkmark$ Positive (c) Non-Negative (d) None of these
- If the initial ray $\overrightarrow{OA}$ rotates in anti-clockwise direction in such a way that it coincides with itself, the angle then formed is:
 (a) 180° (b) 270° (c) 300° (d) $\checkmark 360^\circ$
- One rotation in anti-clock wise direction is equal to:
 (a) 180° (b) 270° (c) $\checkmark 360^\circ$ (d) 90°
- Straight line angle is equal to
 (a) $\frac{1}{2}$ rotation (b) π radian (c) 180° (d) $\checkmark$ All of these
- One right angle is equal to
 (a) $\frac{\pi}{2}$ radian (b) 90° (c) $\frac{1}{4}$ rotation (d) $\checkmark$ All of these
- 1° is equal to
 (a) 30 minutes (b) $\checkmark$ 60 minutes (c) $\frac{1}{60}$ minutes (d) $\frac{1}{2}$ minutes
- 1° is equal to
 (a) $360''$ (b) $\checkmark 3600''$ (c) $(\frac{1}{360})'$ (d) $60''$

10. 60^{th} part of 1° is equal to

- (a) One second (b) ✓ One minute (c) 1 Radian (d) π radian

11. 60^{th} part of $1'$ is equal to

- (a) $1'$ (b) ✓ $1''$ (c) $60''$ (d) $3600''$

12. 3600^{th} part of 1° is equal to

- (a) $1'$ (b) ✓ $1''$ (c) $60''$ (d) $3600''$

13. Sexagesimal system is also called

- (a) German System (b) ✓ English System (c) C.G.S System (d) SI System

14. $16^\circ 30'$ equal to

- (a) ✓ 16.5° (b) $\frac{32^\circ}{2}$ (c) 16.05° (d) 16.2°

15. Conversion of 21.256° to $D^\circ M' S''$ form is:

- (a) $21^\circ 25' 6''$ (b) $21^\circ 40' 27''$ (c) ✓ $21^\circ 15' 22''$ (d) $21^\circ 30' 2''$

16. The angle subtended at the center of a circle by an arc whose length is equal to the radius of the circle is called:

- (a) 1 Degree (b) $1'$ (c) ✓ 1 Radian (d) $1''$

17. The system of angular measurement in which the angle is measured in radian is called:

- (a) Sexagesimal System (b) ✓ Circular System (c) English System (d) Gradient System

18. Relation between the length of arc of a circle and the circular measure of its central angle is:

- (a) $l = \frac{r}{\theta}$ (b) $\theta = lr$ (c) ✓ $\theta = \frac{l}{r}$ (d) $l = \frac{1}{2}r^2\theta$

19. With usual notation, if $l = 6\text{cm}$, $r = 2\text{cm}$, then unit of θ is:

- (a) cm (b) cm^2 (c) ✓ No unit (d) cm^3

20. 1° is equal to:

- (a) $\left(\frac{\pi}{180}\right)^\circ$ (b) $\frac{180}{\pi} \text{ rad}$ (c) $\left(\frac{180}{\pi}\right) \text{ rad}$ (d) ✓ $\frac{\pi}{180} \text{ rad}$

21. 1° is equal to:

- (a) 0.175 rad (b) ✓ 0.0175 rad (c) 1.75 rad (d) 0.00175 rad

22. 1 radian is equal to

- (a) $\left(\frac{\pi}{180}\right)^\circ$ (b) $\frac{180}{\pi} \text{ rad}$ (c) ✓ $\left(\frac{180}{\pi}\right)^\circ$ (d) $\frac{\pi}{180} \text{ rad}$

23. 1 radian is equal to:

- (a) ✓ 57.296° (b) 5.7296° (c) 175.27° (d) 17.5276

24. 3 radian is:

- (a) ✓ 171.888° (b) 120° (c) 300° (d) 270°

25. $105^\circ = \underline{\hspace{2cm}} \text{radian}$

- (a) ✓ $\frac{7\pi}{12}$ (b) $\frac{2\pi}{3}$ (c) $\frac{5\pi}{12}$ (d) $\frac{5\pi}{6}$

26. $3'' = \underline{\hspace{2cm}} \text{radian}$

- (a) $\frac{53\pi}{270}$ (b) ✓ $\frac{\pi}{216000}$ (c) $\frac{41\pi}{720}$ (d) $\frac{27721\pi}{32400}$

27. $\frac{\pi}{4} \text{ radian} = \underline{\hspace{2cm}} \text{degree}$

- (a) ✓ 45° (b) 30° (c) 60° (d) 75°

28. Circular measure of angle between the hands of a watch at 4'O clock is

- (a) 45° (b) ✓ 120° (c) $\frac{3\pi}{2}$ (d) 270°

29. If $l = 1.5 \text{ cm}$ & $r = 2.5 \text{ cm}$ then θ is equal to:

- (a) ✓ $\frac{3}{5}$ (b) $\frac{5}{3}$ (c) 3.75 (d) None

30. If $\theta = 45^\circ$, $r = 18\text{mm}$, then $l =$

- (a) ✓ $\frac{9}{2}\pi$ (b) $\frac{2}{9}\pi$ (c) 812mm (d) 810mm

31. Area of sector of circle of radius r is:

- (a) ✓ $\frac{1}{2}r^2\theta$ (b) $\frac{1}{2}r\theta^2$ (c) $\frac{1}{2}(r\theta)^2$ (d) $\frac{1}{2r^2\theta}$

32. Angles with same initial and terminal sides are called:

- (a) Acute angles (b) Allied Angles (c) ✓ Coterminal angles (d) Quadrantal angles

33. If angle θ is in degree, then the angle coterminal with θ is:

- (a) $\theta + 180^\circ k, k \in \mathbb{Z}$ (b) ✓ $\theta + 360^\circ k, k \in \mathbb{Z}$ (c) $\theta + 90^\circ k, k \in \mathbb{Z}$ (d) $\theta + 60^\circ k, k \in \mathbb{Z}$

34. If angle θ is in degree, then the angle coterminal with θ is:

- (a) ✓ $\theta + 2\pi k, k \in \mathbb{Z}$ (b) $\theta + \pi k, k \in \mathbb{Z}$ (c) $\theta + \frac{\pi}{2}k, k \in \mathbb{Z}$ (d) $\theta + \frac{\pi}{3}k, k \in \mathbb{Z}$

35. An angle is in standard position, if its vertex is

- (a) ✓ At origin (b) at x - axis (c) at y - axis (d) in 1^{st} Quad Only

36. If initial and the terminal side of an angle falls on x – axis or y – axis then it is called:

- (a) Coterminal angle (b) ☒ Quadrantal angle (c) Allied angle (d) None of these

37. $0^\circ, 90^\circ, 180^\circ, 270^\circ$ and 360° are called

- (a) Coterminal angle (b) ☒ Quadrantal angle (c) Allied angle (d) None of these

38. $\sin^2 \theta + \cos^2 \theta$ is equal to:

- (a) 0 (b) -1 (c) 2 (d) ☒ 1

39. $1 + \tan^2 \theta$ is equal to:

- (a) $\csc^2 \theta$ (b) $\sin^2 \theta$ (c) ☒ $\sec^2 \theta$ (d) $\tan^2 \theta$

40. $\csc^2 \theta - \cot^2 \theta$ is equal to:

- (a) 0 (b) ☒ 1 (c) -1 (d) 2

41. If $\sin \theta < 0$ and $\cos \theta > 0$ then the terminal arm of angle lies in Quad.

- (a) I (b) II (c) III (d) ☒ IV

42. If $\cot \theta > 0$ and $\operatorname{cosec} \theta > 0$ then the terminal arm of angle lies in Quad.

- (a) ☒ I (b) II (c) III (d) IV

43. If $\tan \theta < 0$ and $\operatorname{cosec} \theta > 0$ then the terminal arm of angle lies in Quad.

- (a) I (b) ☒ II (c) III (d) IV

44. If $\sec \theta < 0$ and $\sin \theta < 0$ then the terminal arm of angle lies in Quad.

- (a) I (b) II (c) ☒ III (d) IV

45. In right angle triangle, the measure of the side opposite to 30° is:

- (a) ☒ Half of Hypotenuse (b) Half of Base (c) Double of base (d) None of these

46. The point $(0, 1)$ lies on the terminal side of angle:

- (a) 0° (b) ☒ 90° (c) 180° (d) 270°

47. The point $(-1, 0)$ lies on the terminal side of angle:

- (a) 0° (b) 90° (c) ☒ 180° (d) 270°

48. The point $(0, -1)$ lies on the terminal side of angle:

- (a) 0° (b) 90° (c) 180° (d) ☒ 270°

49. $2\sin 45^\circ + \frac{1}{2}\operatorname{Cosec} 45^\circ =$

- (a) $\sqrt{\frac{2}{3}}$ (b) ☒ $\frac{3}{\sqrt{2}}$ (c) -1 (d) 1

50. Domain of $\sin \theta$ is:

- (a) ☒ $\mathbb{R}$ (b) $\theta \in \mathbb{R}$ but $\theta \neq n\pi, n \in \mathbb{Z}$ (c) $\theta \in \mathbb{R}$ but $\theta \neq \frac{(2n+1)\pi}{2}, n \in \mathbb{Z}$ (d) None of these

51. Domain of $\cos \theta$ is:

- (a) ☒ $\mathbb{R}$ (b) $\theta \in \mathbb{R}$ but $\theta \neq n\pi, n \in \mathbb{Z}$ (c) $\theta \in \mathbb{R}$ but $\theta \neq \frac{(2n+1)\pi}{2}, n \in \mathbb{Z}$ (d) None of these

52. Domain of $\tan \theta$ is:

- (a) $\mathbb{R}$ (b) $\theta \in \mathbb{R}$ but $\theta \neq n\pi, n \in \mathbb{Z}$ (c) ☒ $\theta \in \mathbb{R}$ but $\theta \neq \frac{(2n+1)\pi}{2}, n \in \mathbb{Z}$ (d) None of these

53. Domain of $\sec \theta$ is:

- (a) $\mathbb{R}$ (b) $\theta \in \mathbb{R}$ but $\theta \neq n\pi, n \in \mathbb{Z}$ (c) ☒ $\theta \in \mathbb{R}$ but $\theta \neq \frac{(2n+1)\pi}{2}, n \in \mathbb{Z}$ (d) None of these

54. $\operatorname{cosec} \theta \sec \theta \sin \theta \cos \theta =$

- (a) ☒ 1 (b) 0 (c) $\sin \theta$ (d) $\cos \theta$

55. $(\sec \theta + \tan \theta)(\sec \theta - \tan \theta) =$

- (a) ☒ 1 (b) 0 (c) $\sec \theta$ (d) $\tan \theta$

56. $\frac{1 - \sin \theta}{\cos \theta} =$

- (a) $\frac{\cos \theta}{1 - \sin \theta}$ (b) ☒ $\frac{\cos \theta}{1 + \sin \theta}$ (c) $\frac{\sin \theta}{1 - \cos \theta}$ (d) $\frac{\sin \theta}{1 + \cos \theta}$

UNIT # 10 Trigonometric Identities

Each question has four possible answer. Tick the correct answer.

1. Distance between the points $A(3, 8)$ & $B(5, 6)$ is:

- (a) ☒ $2\sqrt{2}$ (b) 3 (c) 4 (d) $\sqrt{2}$

2. Fundamental law of trigonometry is, $\cos(\alpha - \beta)$

- (a) ☒ $\cos \alpha \cos \beta + \sin \alpha \sin \beta$ (b) $\cos \alpha \cos \beta - \sin \alpha \sin \beta$
(c) $\sin \alpha \cos \beta + \cos \alpha \sin \beta$ (d) $\sin \alpha \cos \beta - \cos \alpha \sin \beta$

3. $\cos(\alpha + \beta)$ is equal to:

- (a) $\cos \alpha \cos \beta + \sin \alpha \sin \beta$ (b) ☒ $\cos \alpha \cos \beta - \sin \alpha \sin \beta$
(c) $\sin \alpha \cos \beta + \cos \alpha \sin \beta$ (d) $\sin \alpha \cos \beta - \cos \alpha \sin \beta$

4. $\sin(\alpha + \beta)$ is equal to:

- (a) $\cos\alpha\cos\beta + \sin\alpha\sin\beta$ (b) $\cos\alpha\cos\beta - \sin\alpha\sin\beta$
 (c) ☒ $\sin\alpha\cos\beta + \cos\alpha\sin\beta$ (d) $\sin\alpha\cos\beta - \cos\alpha\sin\beta$

5. $\sin(\alpha - \beta)$ is equal to:

- (a) $\cos\alpha\cos\beta + \sin\alpha\sin\beta$ (b) $\cos\alpha\cos\beta - \sin\alpha\sin\beta$
 (c) $\sin\alpha\cos\beta + \cos\alpha\sin\beta$ (d) ☒ $\sin\alpha\cos\beta - \cos\alpha\sin\beta$

6. $\cos\left(\frac{\pi}{2} - \beta\right) =$

- (a) $\cos\beta$ (b) $-\cos\beta$ (c) ☒ $\sin\beta$ (d) $-\sin\beta$

7. $\cos\left(\beta + \frac{\pi}{2}\right) =$

- (a) $\cos\beta$ (b) $-\cos\beta$ (c) $\sin\beta$ (d) ☒ $-\sin\beta$

8. $\sin\left(\beta - \frac{\pi}{2}\right) =$

- (a) $\cos\beta$ (b) ☒ $-\cos\beta$ (c) $\sin\beta$ (d) ☒ $-\sin\beta$

9. $\cos(2\pi - \theta) =$

- (a) ☒ $\cos\theta$ (b) $-\cos\theta$ (c) $\sin\theta$ (d) $-\sin\theta$

10. $\sin(2\pi - \theta) =$

- (a) $\cos\theta$ (b) $-\cos\theta$ (c) $\sin\theta$ (d) ☒ $-\sin\theta$

11. $\tan(\alpha + \beta) =$

- (a) $\frac{\tan\alpha - \tan\beta}{1 + \tan\alpha\tan\beta}$ (b) ☒ $\frac{\tan\alpha + \tan\beta}{1 - \tan\alpha\tan\beta}$ (c) $\frac{\tan\alpha - \tan\beta}{1 - \tan\alpha\tan\beta}$ (d) $\frac{\tan\alpha + \tan\beta}{1 - \tan\alpha\tan\beta}$

12. $\tan(\alpha - \beta) =$

- (a) ☒ $\frac{\tan\alpha - \tan\beta}{1 + \tan\alpha\tan\beta}$ (b) $\frac{\tan\alpha + \tan\beta}{1 - \tan\alpha\tan\beta}$ (c) $\frac{\tan\alpha - \tan\beta}{1 - \tan\alpha\tan\beta}$ (d) $\frac{\tan\alpha + \tan\beta}{1 - \tan\alpha\tan\beta}$

13. Angles associated with basic angles of measure θ to a right angle or its multiple are called:

- (a) Coterminal angle (b) angle in standard position (c) ☒ Allied angle (d) obtuse angle

14. $\tan\left(\frac{\pi}{2} - \theta\right) =$

- (a) ☒ $\cot\theta$ (b) $\tan\theta$ (c) $-\cot\theta$ (d) $-\tan\theta$

15. $\tan\left(\frac{\pi}{2} + \theta\right) =$

- (a) $\cot\theta$ (b) $\tan\theta$ (c) ☒ $-\cot\theta$ (d) $-\tan\theta$

16. $\sin\left(\frac{3\pi}{2} + \theta\right) =$

- (a) $\sin\theta$ (b) $\cos\theta$ (c) $-\sin\theta$ (d) ☒ $-\cos\theta$

17. $\cos 315^\circ$ is equal to:

- (a) 1 (b) 0 (c) ☒ $\frac{1}{\sqrt{2}}$ (d) $\frac{\sqrt{3}}{2}$

18. $\tan(-135^\circ)$ is equal to:

- (a) ☒ 1 (b) 0 (c) $\frac{1}{\sqrt{3}}$ (d) -1

19. $\sec(-300^\circ) =$

- (a) 1 (b) ☒ 0 (c) 2 (d) -1

20. $\sin(180^\circ + \alpha)\sin(90^\circ - \alpha) =$

- (a) ☒ $\sin\alpha\cos\alpha$ (b) $-\sin\alpha\cos\alpha$ (c) $\cos\gamma$ (d) $-\cos\gamma$

21. If α, β and γ are the angles of a triangle ABC then $\sin(\alpha + \beta) =$

- (a) ☒ $\sin\gamma$ (b) $-\sin\gamma$ (c) $\cos\gamma$ (d) $-\cos\gamma$

22. If α, β and γ are the angles of a triangle ABC then $\cos\left(\frac{\alpha + \beta}{2}\right) =$

- (a) ☒ $\sin\frac{\gamma}{2}$ (b) $-\sin\frac{\gamma}{2}$ (c) $\cos\frac{\gamma}{2}$ (d) $-\cos\frac{\gamma}{2}$

23. If α, β and γ are the angles of a triangle ABC then $\cos(\alpha + \beta) =$

- (a) $\sin\gamma$ (b) $-\sin\gamma$ (c) $\cos\gamma$ (d) ☒ $-\cos\gamma$

24. $\frac{\cos 11^\circ + \sin 11^\circ}{\cos 11^\circ - \sin 11^\circ} =$

- (a) ☒ $\tan 56^\circ$ (b) $\tan 34^\circ$ (c) $\cot 56^\circ$ (d) $\cot 34^\circ$

25. $\sin 2\alpha$ is equal to:

- (a) $\cos^2 \alpha - \sin^2 \alpha$ (b) $1 + \cos 2\alpha$ (c) ☒ $2\sin\alpha\cos\alpha$ (d) $2\sin 2\alpha\cos 2\alpha$

26. $\cos 2\alpha =$

- (a) $\cos^2 \alpha - \sin^2 \alpha$ (b) $1 - 2\sin^2 \alpha$ (c) $2\cos^2 \alpha - 1$ (d) ☒ All of these

27. $\tan 2\alpha =$

- (a) $\frac{2\tan\alpha}{1 + \tan^2 \alpha}$ (b) ☒ $\frac{2\tan\alpha}{1 - \tan^2 \alpha}$ (c) $\frac{2\tan^2 \alpha}{1 - \tan^2 \alpha}$ (d) $\frac{\tan^2 \alpha}{1 - \tan^2 \alpha}$

28. $\sin\alpha + \sin\beta$ is equal to:

- (a) ☒ $2\sin\left(\frac{\alpha + \beta}{2}\right)\cos\left(\frac{\alpha - \beta}{2}\right)$ (b) $2\cos\left(\frac{\alpha + \beta}{2}\right)\sin\left(\frac{\alpha - \beta}{2}\right)$
 (c) $-2\sin\left(\frac{\alpha + \beta}{2}\right)\sin\left(\frac{\alpha - \beta}{2}\right)$ (d) $2\cos\left(\frac{\alpha + \beta}{2}\right)\cos\left(\frac{\alpha - \beta}{2}\right)$

29. $\sin\alpha - \sin\beta$ is equal to:

- (b) $2 \sin\left(\frac{\alpha+\beta}{2}\right) \cos\left(\frac{\alpha-\beta}{2}\right)$ (b) ✓ $2 \cos\left(\frac{\alpha+\beta}{2}\right) \sin\left(\frac{\alpha-\beta}{2}\right)$
 (c) $-2 \sin\left(\frac{\alpha+\beta}{2}\right) \sin\left(\frac{\alpha-\beta}{2}\right)$ (d) $2 \cos\left(\frac{\alpha+\beta}{2}\right) \cos\left(\frac{\alpha-\beta}{2}\right)$

30. $\cos\alpha + \cos\beta$ is equal to:

- (c) $2 \sin\left(\frac{\alpha+\beta}{2}\right) \cos\left(\frac{\alpha-\beta}{2}\right)$ (b) $2 \cos\left(\frac{\alpha+\beta}{2}\right) \sin\left(\frac{\alpha-\beta}{2}\right)$
 (c) $-2 \sin\left(\frac{\alpha+\beta}{2}\right) \sin\left(\frac{\alpha-\beta}{2}\right)$ (d) ✓ $2 \cos\left(\frac{\alpha+\beta}{2}\right) \cos\left(\frac{\alpha-\beta}{2}\right)$

31. $\cos\alpha - \cos\beta$ is equal to:

- (d) $2 \sin\left(\frac{\alpha+\beta}{2}\right) \cos\left(\frac{\alpha-\beta}{2}\right)$ (b) $2 \cos\left(\frac{\alpha+\beta}{2}\right) \sin\left(\frac{\alpha-\beta}{2}\right)$
 (c) ✓ $-2 \sin\left(\frac{\alpha+\beta}{2}\right) \sin\left(\frac{\alpha-\beta}{2}\right)$ (d) $2 \cos\left(\frac{\alpha+\beta}{2}\right) \cos\left(\frac{\alpha-\beta}{2}\right)$

32. Which is the allied angle

- (a) ✓ $90^\circ + \theta$ (b) $60^\circ + \theta$ (c) $45^\circ + \theta$ (d) $30^\circ + \theta$

33. $2\sin 7\theta \cos 3\theta =$

- (a) ✓ $\sin 10\theta + \sin 4\theta$ (b) $\sin 5\theta - \sin 2\theta$ (c) $\cos 10\theta + \cos 4\theta$ (d) $\cos 5\theta - \cos 2\theta$

34. $2\cos 5\theta \sin 3\theta =$

- (b) ✓ $\sin 8\theta - \sin 2\theta$ (b) $\sin 8\theta + \sin 2\theta$ (c) $\cos 8\theta + \cos 2\theta$ (d) $\cos 8\theta - \cos 2\theta$

UNIT # 11 Trigonometric Functions and their Graphs

Each question has four possible answer. Tick the correct answer.

1. Domain of $y = \sin x$ is

- (a) ✓ $-\infty < x < \infty$ (b) $-1 \leq x \leq 1$ (c) $-\infty < x < \infty, x \neq n\pi, n \in \mathbb{Z}$ (d) $x \geq 1, x \leq -1$

2. Domain of $y = \cos x$ is

- (a) ✓ $-\infty < x < \infty$ (b) $-1 \leq x \leq 1$ (c) $-\infty < x < \infty, x \neq n\pi, n \in \mathbb{Z}$ (d) $x \geq 1, x \leq -1$

3. Domain of $y = \tan x$ is

- (a) $-\infty < x < \infty$ (b) $-1 \leq x \leq 1$ (c) ✓ $-\infty < x < \infty, x \neq \frac{2n+1}{2}\pi, n \in \mathbb{Z}$ (d) $x \geq 1, x \leq -1$

4. Domain of $y = \sec x$ is

- (a) $-\infty < x < \infty$ (b) $-1 \leq x \leq 1$ (c) ✓ $-\infty < x < \infty, x \neq \frac{2n+1}{2}\pi, n \in \mathbb{Z}$ (d) $x \geq 1, x \leq -1$

5. Domain of $y = \csc x$ is

- (a) $-\infty < x < \infty$ (b) ✓ $-1 \leq x \leq 1$ (c) $-\infty < x < \infty, x \neq \frac{2n+1}{2}\pi, n \in \mathbb{Z}$ (d) $x \geq 1, x \leq -1$

6. Domain of $y = \cot x$ is

- (a) $-\infty < x < \infty$ (b) ✓ $-1 \leq x \leq 1$ (c) $-\infty < x < \infty, x \neq \frac{2n+1}{2}\pi, n \in \mathbb{Z}$ (d) $x \geq 1, x \leq -1$

7. Range of $y = \sin x$ is

- (a) $\mathbb{R}$ (b) ✓ $-1 \leq y \leq 1$ (c) $(-\infty, 1) \cup (1, \infty)$ (d) $-1 < y < 1$

8. Range of $y = \cos x$ is

- (a) $\mathbb{R}$ (b) ✓ $-1 \leq y \leq 1$ (c) $(-\infty, 1) \cup (1, \infty)$ (d) $-1 < y < 1$

9. Range of $y = \tan x$ is

- (a) ✓ $\mathbb{R}$ (b) $-1 \leq y \leq 1$ (c) $\mathbb{Q}$ (d) $\mathbb{R} - \{0\}$

10. Range of $y = \cot x$ is

- (a) ✓ $\mathbb{R}$ (b) $\mathbb{R} - [-1, 1]$ (c) $\mathbb{R} - \{0\}$ (d) $\mathbb{Z}$

11. Range of $y = \sec x$ is

- (a) $\mathbb{R}$ (b) ✓ $y \geq 1 \text{ or } y \leq -1$ (c) $-1 \leq y \leq 1$ (d) $\mathbb{R} - [-1, 1]$

12. Range of $y = \csc x$ is

- (a) $\mathbb{R}$ (b) ✓ $y \geq 1 \text{ or } y \leq -1$ (c) $-1 \leq y \leq 1$ (d) $\mathbb{R} - [-1, 1]$

13. Smallest +ive number which when added to the original circular measure of the angle gives the same value of the function is called:

- (a) Domain (b) Range (c) Co domain (d) ✓ Period

14. Period of $\sin \theta$ is

- (a) π (b) ✓ 2π (c) -2π (d) $\frac{\pi}{2}$

15. Period of $\csc \theta$ is

- (a) π (b) ✓ 2π (c) -2π (d) $\frac{\pi}{2}$

16. Period of $\cos\theta$ is

- (a) π (b) ☒ 2π (c) -2π (d) $\frac{\pi}{2}$

17. Period of $\tan\theta$ is

- (a) ☒ π (b) 2π (c) -2π (d) $\frac{\pi}{2}$

18. Period of $\cot\theta$ is

- (a) ☒ π (b) 2π (c) -2π (d) $\frac{\pi}{2}$

19. Period of $\sec\theta$ is

- (a) π (b) ☒ 2π (c) -2π (d) $\frac{\pi}{2}$

20. Period of $\tan 4x$ is

- (a) π (b) 2π (c) -2π (d) ☒ $\frac{\pi}{4}$

21. Period of $\cot 3x$ is

- (a) π (b) ☒ $\frac{\pi}{3}$ (c) -2π (d) $\frac{\pi}{4}$

22. Period of $3\cos\frac{x}{5}$ is

- (a) 2π (b) $\frac{\pi}{2}$ (c) π (d) ☒ 10π

23. The graph of trigonometric functions have:

- (a) Break segments (b) Sharp corners (c) Straight line segments (d) ☒ smooth curves

24. Curves of the trigonometric functions repeat after fixed intervals because trigonometric functions are

- (a) Simple (b) linear (c) quadratic (d) ☒ periodic

25. The graph of $y = \cos x$ lies between the horizontal line $y = -1$ and

- (a) ☒ $+1$ (b) 0 (c) 2 (d) -2

UNIT # 12 Application of Trigonometry

Each question has four possible answer. Tick the correct answer.

1. A "Triangle" has :

- (a) Two elements (b) 3 elements (c) 4 elements (d) ☒ 6 elements

2. When we look an object above the horizontal ray, the angle formed is called angle of:

- (a) ☒ Elevation (b) depression (c) incidence (d) reflects

3. When we look an object below the horizontal ray, the angle formed is called angle of:

- (a) Elevation (b) ☒ depression (c) incidence (d) reflects

4. A triangle which is not right is called:

- (a) ☒ Oblique triangle (b) Isosceles triangle (c) Scalene triangle (d) Right isosceles triangle

5. To solve an oblique triangle we use:

- (a) ☒ Law of Sine (b) Law of Cosine (c) Law of Tangents (d) All of these

6. In any triangle ABC , $\frac{b^2+c^2-a^2}{2bc} =$

- (a) ☒ $\cos\alpha$ (b) $\sin\alpha$ (c) $\cos\beta$ (d) $\cos\gamma$

7. Which can be reduced to Pythagoras theorem,

- (a) Law of sine (b) ☒ law of cosine (c) law of tangents (d) Half angle formulas

8. In any triangle ABC , if $\beta = 90^\circ$, then $b^2 = a^2 + c^2 - 2accos\beta$ becomes:

- (a) Law of sine (b) Law of tangents (c) ☒ Law of cosine (d) None of these

9. In any triangle ABC , law of tangent is :

- (a) $\frac{a-b}{a+b} = \frac{\tan(\alpha-\beta)}{\tan(\alpha+\beta)}$ (b) $\frac{a+b}{a-b} = \frac{\tan(\alpha+\beta)}{\tan(\alpha-\beta)}$ (c) ☒ $\frac{a-b}{a+b} = \frac{\tan\frac{\alpha-\beta}{2}}{\tan\frac{\alpha+\beta}{2}}$ (d) $\frac{a-b}{a+b} = \frac{\tan\frac{\alpha+\beta}{2}}{\tan\frac{\alpha-\beta}{2}}$

10. In any triangle ABC , $\sqrt{\frac{(s-a)(s-b)}{ab}} =$

- (a) $\sin\frac{\alpha}{2}$ (b) $\sin\frac{\beta}{2}$ (c) ☒ $\sin\frac{\gamma}{2}$ (d) $\cos\frac{\alpha}{2}$

11. In any triangle ABC , $\sqrt{\frac{(s-b)(s-c)}{bc}} =$

- (a) ☒ $\sin\frac{\alpha}{2}$ (b) $\sin\frac{\beta}{2}$ (c) $\sin\frac{\gamma}{2}$ (d) $\cos\frac{\alpha}{2}$

12. In any triangle ABC , $\sqrt{\frac{(s-a)(s-c)}{ac}} =$

- (a) $\sin\frac{\alpha}{2}$ (b) ☒ $\sin\frac{\beta}{2}$ (c) $\sin\frac{\gamma}{2}$ (d) $\cos\frac{\alpha}{2}$

13. In any triangle ABC , $\cos\frac{\alpha}{2} =$

- (a) $\sqrt{\frac{s(s-a)}{ab}}$ (b) $\sqrt{\frac{s(s-b)}{ac}}$ (c) ☒ $\sqrt{\frac{s(s-a)}{bc}}$ (d) $\sqrt{\frac{s(s-c)}{ab}}$

14. In any triangle ABC , $\cos \frac{\beta}{2} =$

- (a) $\sqrt{\frac{s(s-a)}{ab}}$ (b) ☒ $\sqrt{\frac{s(s-b)}{ac}}$ (c) $\sqrt{\frac{s(s-a)}{bc}}$ (d) $\sqrt{\frac{s(s-c)}{ab}}$

15. In any triangle ABC , $\cos \frac{\gamma}{2} =$

- (a) $\sqrt{\frac{s(s-a)}{ab}}$ (b) $\sqrt{\frac{s(s-b)}{ac}}$ (c) $\sqrt{\frac{s(s-a)}{bc}}$ (d) ☒ $\sqrt{\frac{s(s-c)}{ab}}$

16. In any triangle ABC , with usual notations, s is equal to

- (a) $a + b + c$ (b) $\frac{a+b+c}{3}$ (c) ☒ $\frac{a+b+c}{2}$ (d) $\frac{abc}{2}$

17. In any triangle ABC , $\sqrt{\frac{s(s-c)}{(s-a)(s-b)}} =$

- (a) $\sin \frac{\gamma}{2}$ (b) $\cos \frac{\gamma}{2}$ (c) $\tan \frac{\gamma}{2}$ (d) ☒ $\cot \frac{\gamma}{2}$

18. In any triangle ABC , $\sqrt{\frac{(s-a)(s-b)}{s(s-c)}} =$

- (a) $\sin \frac{\gamma}{2}$ (b) $\cos \frac{\gamma}{2}$ (c) ☒ $\tan \frac{\gamma}{2}$ (d) $\cot \frac{\gamma}{2}$

19. To solve an oblique triangles when measure of three sides are given, we can use:

- (a) ☒ Hero's formula (b) Law of cosine (c) Law of sine (d) Law of tangents

20. The smallest angle of $\triangle ABC$, when $a = 37.34$, $b = 3.24$, $c = 35.06$ is

- (a) α (b) ☒ β (c) γ (d) cannot be determined

21. In any triangle ABC Area if triangle is :

- (a) $bc \sin \alpha$ (b) $\frac{1}{2} ca \sin \alpha$ (c) $\frac{1}{2} ab \sin \beta$ (d) ☒ $\frac{1}{2} abc \sin \gamma$

22. The circle passing through the three vertices of a triangle is called:

- (a) ☒ Circum circle (b) in-circle (c) ex-centre (d) escribed circle

23. The point of intersection of the right bisectors of the sides of the triangle is :

- (a) ☒ Circum centre (b) In-centre (c) Escribed center (d) Diameter

24. In any triangle ABC , with usual notations, $\frac{a}{2 \sin \alpha} =$

- (a) r (b) r_1 (c) ☒ R (d) Δ

25. In any triangle ABC , with usual notations, $\frac{a}{\sin \beta} =$

- (a) $2r$ (b) $2r_1$ (c) ☒ $2R$ (d) 2Δ

26. In any triangle ABC , with usual notations, $\sin \gamma =$

- (a) R (b) ☒ $\frac{c}{2R}$ (c) $\frac{2R}{c}$ (d) $\frac{R}{2}$

27. In any triangle ABC , with usual notations, $abc =$

- (a) R (b) R_s (c) ☒ $4R\Delta$ (d) $\frac{\Delta}{s}$

28. In any triangle ABC , with usual notations, $\frac{\Delta}{s-a} =$

- (a) r (b) R (c) ☒ r_1 (d) r_2

29. In any triangle ABC , with usual notations, $\frac{\Delta}{s-b} =$

- (a) r (b) R (c) r_1 (d) ☒ r_2

30. In any triangle ABC , with usual notations, $\frac{\Delta}{s-c} =$

- (a) ☒ r_3 (b) R (c) r_1 (d) r_2

31. In any triangle ABC , with usual notation, $r:R:r_1 =$

- (a) $3:2:1$ (b) $1:2:2$ (c) ☒ $1:2:3$ (d) $1:1:1$

32. In any triangle ABC , with usual notation, $r:R:r_1:r_2:r_3 =$

- (a) $3:3:3:2:1$ (b) $1:2:2:3:3$ (c) ☒ $1:2:3:3:3$ (d) $1:1:1:1:1$

33. In a triangle ABC , if $\beta = 60^\circ$, $\gamma = 15^\circ$ then $\alpha =$

- (a) 90° (b) 180° (c) 150° (d) ☒ 105°

UNIT # 13 Inverse trigonometric functions

Each question has four possible answer. Tick the correct answer.

1. If $y = \sin x$, then Domain is :

- (a) ☒ $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) $0 \leq x \leq \pi$ (c) $[0, \pi], x \neq \frac{\pi}{2}$ (d) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right], x \neq 0$

2. If $y = \cos x$, then Domain is :

- (a) $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) ☒ $0 \leq x \leq \pi$ (c) $[0, \pi], x \neq \frac{\pi}{2}$ (d) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right], x \neq 0$

3. If $y = \sec x$, then Domain is :

- (a) $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) $0 \leq x \leq \pi$ (c) ☒ $[0, \pi], x \neq \frac{\pi}{2}$ (d) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right], x \neq 0$

4. If $y = \operatorname{Cosec} x$, then Domain is :

- (a) $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) $0 \leq x \leq \pi$ (c) $[0, \pi], x \neq \frac{\pi}{2}$ (d) ☒ $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right], x \neq 0$

5. If $y = \tan x$, then Domain is :

- (a) $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) ☒ $0 \leq x \leq \pi$ (c) $[0, \pi], x \neq \frac{\pi}{2}$ (d) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right], x \neq 0$

6. If $y = \cot x$, then Domain is :

- (a) $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) $0 \leq x \leq \pi$ (c) ☒ $[0, \pi], x \neq \frac{\pi}{2}$ (d) $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right], x \neq 0$

7. If $y = \sin x$, then range is :

- (a) ☒ $-1 \leq x \leq 1$ (b) $(-\infty, +\infty)$ or R (c) $y \leq -1$ or $y \geq 1$ (d) $y < -1$ or $y > 1$

8. If $y = \cos x$, then range is :

- (a) ☒ $-1 \leq x \leq 1$ (b) $(-\infty, +\infty)$ or R (c) $y \leq -1$ or $y \geq 1$ (d) $y < -1$ or $y > 1$

9. If $y = \tan x$, then range is :

- (a) $-1 \leq x \leq 1$ (b) ☒ $(-\infty, +\infty)$ or R (c) $y \leq -1$ or $y \geq 1$ (d) $y < -1$ or $y > 1$

10. If $y = \cot x$, then range is :

- (a) $-1 \leq x \leq 1$ (b) ☒ $(-\infty, +\infty)$ or R (c) $y \leq -1$ or $y \geq 1$ (d) $y < -1$ or $y > 1$

11. If $y = \operatorname{Cosec} x$, then range is :

- (a) $-1 \leq x \leq 1$ (b) $(-\infty, +\infty)$ or R (c) ☒ $y \leq -1$ or $y \geq 1$ (d) $y < -1$ or $y > 1$

12. If $y = \sec x$, then range is :

- (a) $-1 \leq x \leq 1$ (b) $(-\infty, +\infty)$ or R (c) ☒ $y \leq -1$ or $y \geq 1$ (d) $y < -1$ or $y > 1$

13. If $y = \sin^{-1} x$, then domain is:

- (a) ☒ $-1 \leq x \leq 1$ (b) $(-\infty, +\infty)$ or R (c) $x \geq -1$ or $x \leq 1$ (d) $x \leq -1$ or $x \geq 1$

14. If $y = \cos^{-1} x$, then domain is:

- (a) ☒ $-1 \leq x \leq 1$ (b) $(-\infty, +\infty)$ or R (c) $x \geq -1$ or $x \leq 1$ (d) $x \leq -1$ or $x \geq 1$

15. If $y = \tan^{-1} x$, then domain is:

- (a) $-1 \leq x \leq 1$ (b) ☒ $(-\infty, +\infty)$ or R (c) $x \geq -1$ or $x \leq 1$ (d) $x \leq -1$ or $x \geq 1$

16. If $y = \cot^{-1} x$, then domain is:

- (a) $-1 \leq x \leq 1$ (b) ☒ $(-\infty, +\infty)$ or R (c) $x \geq -1$ or $x \leq 1$ (d) $x \leq -1$ or $x \geq 1$

17. If $y = \sec^{-1} x$, then domain is:

- (a) $-1 \leq x \leq 1$ (b) $(-\infty, +\infty)$ or R (c) ☒ $x \geq -1$ or $x \leq 1$ (d) $x \leq -1$ or $x \geq 1$

18. If $y = \operatorname{Cosec}^{-1} x$, then domain is:

- (a) $-1 \leq x \leq 1$ (b) $(-\infty, +\infty)$ or R (c) ☒ $x \geq -1$ or $x \leq 1$ (d) $x \leq -1$ or $x \geq 1$

19. If $y = \sin^{-1} x$, then range is:

- (a) ☒ $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) $0 \leq x \leq \pi$ (c) $-\frac{\pi}{2} < x < \frac{\pi}{2}$ (d) $0 < x < \pi$

20. If $y = \cos^{-1} x$, then range is:

- (a) $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) ☒ $0 \leq x \leq \pi$ (c) $-\frac{\pi}{2} < x < \frac{\pi}{2}$ (d) $0 < x < \pi$

21. If $y = \tan^{-1} x$, then range is:

- (a) $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) $0 \leq x \leq \pi$ (c) ☒ $-\frac{\pi}{2} < x < \frac{\pi}{2}$ (d) $0 < x < \pi$

22. If $y = \cot^{-1} x$, then range is:

- (a) $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) $0 \leq x \leq \pi$ (c) $-\frac{\pi}{2} < x < \frac{\pi}{2}$ (d) ☒ $0 < x < \pi$

23. If $y = \sec^{-1} x$, then range is:

- (a) ☒ $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) $0 \leq x \leq \pi$ (c) $-\frac{\pi}{2} < x < \frac{\pi}{2}$ (d) $0 < x < \pi$

24. If $y = \operatorname{Cosec}^{-1} x$, then range is:

- (a) $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ (b) ☒ $0 \leq x \leq \pi$ (c) $-\frac{\pi}{2} < x < \frac{\pi}{2}$ (d) $0 < x < \pi$

25. Inverse of a function exist only if it is:

- (a) Trigonometric function (b) ☒ $(1-1)$ function (c) onto function (d) an into function

26. $\sin^{-1} x =$

- (a) ☒ $\frac{\pi}{2} - \cos^{-1} x$ (b) $\frac{\pi}{2} - \sin^{-1} x$ (c) $\frac{\pi}{2} + \cos^{-1} x$ (d) $\frac{\pi}{2} - \operatorname{cosec}^{-1} x$

27. $\cos^{-1} x =$

- (a) $\frac{\pi}{2} - \cos^{-1} x$ (b) ☒ $\frac{\pi}{2} - \sin^{-1} x$ (c) $\frac{\pi}{2} + \cos^{-1} x$ (d) $\frac{\pi}{2} - \operatorname{cosec}^{-1} x$

28. $\cos^{-1} x =$

- (a) ☒ $\frac{\pi}{2} - \sec^{-1} x$ (b) $\frac{\pi}{2} - \sin^{-1} x$ (c) $\frac{\pi}{2} + \sec^{-1} x$ (d) $\frac{\pi}{2} - \operatorname{cosec}^{-1} x$

29. $\sec^{-1} x =$

- (a) $\frac{\pi}{2} - \sec^{-1} x$ (b) $\frac{\pi}{2} - \sin^{-1} x$ (c) $\frac{\pi}{2} + \sec^{-1} x$ (d) ☒ $\frac{\pi}{2} - \operatorname{cosec}^{-1} x$

30. $\tan^{-1} x =$

- (a) $\frac{\pi}{2} - \sec^{-1} x$ (b) $\frac{\pi}{2} - \sin^{-1} x$ (c) ☒ $\frac{\pi}{2} - \cot^{-1} x$ (d) $\frac{\pi}{2} - \operatorname{cosec}^{-1} x$

31. $\cot^{-1} x =$

- (a) $\frac{\pi}{2} - \sec^{-1} x$ (b) ☒ $\frac{\pi}{2} - \tan^{-1} x$ (c) $\frac{\pi}{2} + \sec^{-1} x$ (d) $\frac{\pi}{2} - \operatorname{cosec}^{-1} x$

32. $\sin\left(\cos^{-1}\frac{\sqrt{3}}{2}\right) =$

- (a) $\frac{\pi}{6}$ (b) ☒ $\frac{1}{2}$ (c) $-\frac{1}{2}$ (d) $\frac{\sqrt{3}}{2}$

33. $\tan^{-1}(\sqrt{3}) =$

- (a) $\frac{\pi}{6}$ (b) $-\frac{\pi}{6}$ (c) $-\frac{\pi}{3}$ (d) ☒ $\frac{\pi}{3}$

34. $\sin\left(\sin^{-1}\frac{1}{2}\right) =$

- (a) ☒ $\frac{1}{2}$ (b) $\frac{2}{3}$ (c) 2 (d) $\frac{1}{3}$

35. $\sin^{-1}A + \sin^{-1}B =$

- (a) ☒ $\sin^{-1}\left(A\sqrt{1-B^2} + B\sqrt{1-A^2}\right)$ (b) $\sin^{-1}\left(A\sqrt{1-A^2} - B\sqrt{1-B^2}\right)$
(c) $\sin^{-1}\left(B\sqrt{1-A^2} + A\sqrt{1-B^2}\right)$ (d) $\sin^{-1}\left(AB\sqrt{(1-A^2)(1-B^2)}\right)$

36. $\sin^{-1}A - \sin^{-1}B =$

- (a) $\sin^{-1}\left(A\sqrt{1-B^2} + B\sqrt{1-A^2}\right)$ (b) ☒ $\sin^{-1}\left(A\sqrt{1-A^2} - B\sqrt{1-B^2}\right)$
(c) $\sin^{-1}\left(B\sqrt{1-A^2} + A\sqrt{1-B^2}\right)$ (d) $\sin^{-1}\left(AB\sqrt{(1-A^2)(1-B^2)}\right)$

37. $\cos^{-1}A + \cos^{-1}B =$

- (a) $\cos^{-1}\left(AB - \sqrt{(1-A^2)(1-B^2)}\right)$ (b) $\cos^{-1}\left(AB + \sqrt{(1-A^2)(1-B^2)}\right)$
(c) $\cos^{-1}\left(AB - \sqrt{(1+A^2)(1+B^2)}\right)$ (d) $\cos^{-1}\left(AB + \sqrt{(1+A^2)(1+B^2)}\right)$

38. $\cos^{-1}A + \cos^{-1}B =$

- (a) ☒ $\cos^{-1}\left(AB - \sqrt{(1-A^2)(1-B^2)}\right)$ (b) ☒ $\cos^{-1}\left(AB + \sqrt{(1-A^2)(1-B^2)}\right)$
(c) $\cos^{-1}\left(AB - \sqrt{(1+A^2)(1+B^2)}\right)$ (d) $\cos^{-1}\left(AB + \sqrt{(1+A^2)(1+B^2)}\right)$

39. $\tan^{-1}A + \tan^{-1}B =$

- (a) ☒ $\tan^{-1}\left(\frac{A+B}{1-AB}\right)$ (b) $\tan^{-1}\left(\frac{A+B}{1+AB}\right)$ (c) $\tan^{-1}\left(\frac{A-B}{1-AB}\right)$ (d) $\tan^{-1}\left(\frac{A+B}{1+AB}\right)$

40. $\tan^{-1}A + \tan^{-1}B =$

- (a) $\tan^{-1}\left(\frac{A-B}{1+AB}\right)$ (b) ☒ $\tan^{-1}\left(\frac{A+B}{1+AB}\right)$ (c) $\tan^{-1}\left(\frac{A-B}{1-AB}\right)$ (d) $\tan^{-1}\left(\frac{A+B}{1+AB}\right)$

41. $\sin^{-1}(-x) =$

- (a) ☒ $-\sin^{-1}x$ (b) $\sin^{-1}x$ (c) $\pi - \sin^{-1}x$ (d) $\pi - \sin x$

42. $\cos^{-1}(-x) =$

- (a) $-\cos^{-1}x$ (b) $\cos^{-1}x$ (c) ☒ $\pi - \cos^{-1}x$ (d) $\pi - \cos x$

43. $\tan^{-1}(-x) =$

- (a) ☒ $-\tan^{-1}x$ (b) $\tan^{-1}x$ (c) $\pi - \tan^{-1}x$ (d) $\pi - \tan x$

44. $\operatorname{cosec}^{-1}(-x) =$

- (a) ☒ $-\operatorname{cosec}^{-1}x$ (b) $\operatorname{cosec}^{-1}x$ (c) $\pi - \operatorname{cosec}^{-1}x$ (d) $\pi - \operatorname{cosec} x$

45. $\sec^{-1}(-x) =$

- (a) $-\sec^{-1}x$ (b) $\sec^{-1}x$ (c) ☒ $\pi - \sec^{-1}x$ (d) $\pi - \sec x$

46. $\cot^{-1}(-x) =$

- (a) $-\cot^{-1}x$ (b) $\cot^{-1}x$ (c) ☒ $\pi - \cot^{-1}x$ (d) $\pi - \cot x$

UNIT # 14 Solution of Trigonometric Equations

Each question has four possible answer. Tick the correct answer.

1. An equation containing at least one trigonometric function is called:

- (a) Trigonometric function (b) ☒ Trigonometric equation (c) Trigonometric value (d) None

2. If $\sin x = \frac{1}{2}$, then solution in the interval $[0, 2\pi]$ is:

- (a) ☒ $\left\{\frac{\pi}{6}, \frac{5\pi}{6}\right\}$ (b) $\left\{\frac{\pi}{6}, \frac{7\pi}{6}\right\}$ (c) $\left\{\frac{\pi}{3}, \frac{4\pi}{6}\right\}$ (d) $\left\{\frac{\pi}{3}, \frac{2\pi}{6}\right\}$

3. If $\cos x = \frac{1}{2}$, then the reference angle is:

- (a) ☒ $\frac{\pi}{3}$ (b) $-\frac{\pi}{3}$ (c) $\frac{\pi}{6}$ (d) $-\frac{\pi}{6}$

4. If $\sin x = \frac{1}{2}$, then the reference angle is:

- (a) $\frac{\pi}{3}$ (b) $-\frac{\pi}{3}$ (c) ☒ $\frac{\pi}{6}$ (d) $-\frac{\pi}{6}$

5. General solution of $\tan x = 1$ is:

- (a) ☒ $\{\frac{\pi}{4} + n\pi, \frac{5\pi}{4} + n\pi\}$ (b) $\{\frac{\pi}{4} + 2n\pi, \frac{5\pi}{4} + 2n\pi\}$ (c) $\{\frac{\pi}{4} + n\pi, \frac{3\pi}{4} + n\pi\}$ (d) $\{\frac{\pi}{4} + 2n\pi, \frac{3\pi}{4} + 2n\pi\}$

6. If $\tan 2x = -1$, then solution in the interval $[0, \pi]$ is:

- (a) ☒ $\frac{\pi}{8}$ (b) $\frac{\pi}{4}$ (c) $\frac{3\pi}{8}$ (d) $\frac{3\pi}{4}$

7. If $\sin x + \cos x = 0$ then value of $x \in [0, 2\pi]$

- (a) $\{\frac{\pi}{4}, \frac{3\pi}{4}\}$ (b) $\{\frac{\pi}{4}, \frac{7\pi}{4}\}$ (c) ☒ $\{\frac{3\pi}{4}, \frac{7\pi}{4}\}$ (d) $\{\frac{\pi}{4}, \frac{-\pi}{4}\}$

8. General solution of $4\sin x - 8 = 0$ is:

- (a) $\{\pi + 2n\pi\}$ (b) $\{\pi + n\pi\}$ (c) $\{-\pi + n\pi\}$ (d) ☒ not possible

9. General solution of $1 + \cos x = 0$ is:

- (a) ☒ $\{\pi + 2n\pi\}$ (b) $\{\pi + n\pi\}$ (c) $\{-\pi + n\pi\}$ (d) not possible

10. For the general solution, we first find the solution in the interval whose length is equal to its:

- (a) Range (b) domain (c) co-domain (d) ☒ period

11. All trigonometric functions are functions.

- (a) ☒ Periodic (b) continues (c) injective (d) bijective

12. General solution of every trigonometric equation consists of :

- (a) One solution only (b) two solutions (c) ☒ infinitely many solutions (d) no real solution

13. Solution of the equation $2\sin x + \sqrt{3} = 0$ in the 4th quadrant is:

- (a) $\frac{\pi}{2}$ (b) ☒ $\frac{-\pi}{3}$ (c) $\frac{-\pi}{6}$ (d) $\frac{11\pi}{6}$

14. If $\sin x = \cos x$, then general solution is:

- (a) $\{\frac{\pi}{4} + n\pi, n \in \mathbb{Z}\}$ (b) $\{\frac{\pi}{4} + 2n\pi, n \in \mathbb{Z}\}$
(c) ☒ $\{\frac{\pi}{4} + n\pi, \frac{5\pi}{4} + n\pi\}$ (d) $\{\frac{\pi}{4} + n\pi, \frac{5\pi}{4} + n\pi\}$

15. In which quadrant is the solution of the equation $\sin x + 1 = 0$

- (a) 1st and 2nd (b) 2nd and 3rd (c) ☒ 3rd and 4th (d) Only 1st

16. If $\sin x = 0$ then $x =$

- (a) ☒ $n\pi, n \in \mathbb{Z}$ (b) $\frac{n\pi}{2}, n \in \mathbb{Z}$ (c) 0 (d) $\frac{\pi}{2}$

17. If $\tan x = 0$ then $x =$

- (a) ☒ $n\pi, n \in \mathbb{Z}$ (b) $\frac{n\pi}{2}, n \in \mathbb{Z}$ (c) 0 (d) $\frac{\pi}{2}$

18. The solution of the $\cot x = \frac{1}{\sqrt{3}}$ in $[0, \pi]$ is

- (a) $\frac{\pi}{6}$ (b) $\frac{\pi}{4}$ (c) $\frac{2\pi}{3}$ (d) ☒ $\frac{\pi}{3}$

19. One solution of $\sec x = -2$ is :

- (a) $\frac{\pi}{6}$ (b) $\frac{\pi}{4}$ (c) ☒ $\frac{2\pi}{3}$ (d) $\frac{\pi}{3}$

20. $\sin 2x = \frac{\sqrt{3}}{2}$ has two values of x in the interval:

- (a) $[0, \frac{\pi}{2}]$ (b) $[0, 2\pi]$ (c) $[-\pi, \frac{\pi}{2}]$ (d) $[-\frac{\pi}{2}, 0]$

←-----THE END-----→

WITH BEST WISHES BY:-



MUHAMMAD SALMAN

SHERAZI M.Sc(Math) , B.ed