

CHAPTER : 1

1. $\frac{e^{2x}-1}{2e^{2x}}$ is equal to:
- a. $\sin x$
 - b. $\cos x$
 - c. $\sinh x$
 - d. $\cosh x$
2. $\lim_{\theta \rightarrow 0} \frac{1-\cos p\theta}{1+\cos q\theta}$ equals to:
- a. **0**
 - b. $\frac{p}{q}$
 - c. $\frac{p^2}{q^2}$
 - d. $\frac{q^2}{p^2}$
3. $x=3 \cos t, y=3 \sin t$ represent:
- a. line
 - b. circle
 - c. parabola
 - d. hyperbola
4. $\lim_{x \rightarrow 0} \frac{\sin 3x}{\sin 2x}$ is equal to:
- a. $\frac{2}{3}$
 - b. $\frac{3}{2}$
 - c. $\frac{1}{6}$
 - d. $\frac{1}{4}$
5. If $(x) = x^2 - x$ then $f(-2)$ is equal to:
- a. 2
 - b. 6
 - c. 0
 - d. -6
6. If $\lim_{x \rightarrow \infty} \frac{f(x)-1}{x} = \ln a ; a > 0$
- a. a^{-x}
 - b. a^x
 - c. e^{-x}
 - d. e^x
7. $\lim_{x \rightarrow \infty} (1 + \frac{x}{2})^{\frac{1}{x}}$ equals:
- a. e
 - b. e^{-1}
 - c. e^2
 - d. \sqrt{e}
8. $\frac{d}{dx} e^{f(x)}$ equals:
- a. $e^{f'(x)}$
 - b. $e^{f(x)} \cdot f'(x)$
 - c. $\frac{f'(x)}{e^x}$
 - d. $\frac{e^{f(x)}}{f'(x)}$

9. $\lim_{x \rightarrow 2} \frac{x^5 - 32}{x - 2}$ equals:

- a. 40
- c. 80

- b. 60
- d. 120

10. $\lim_{\theta \rightarrow 0} \frac{\sin 7\theta}{\theta}$ is equal to:

- a. $\frac{1}{7}$
- c. 1

- b. 7
- d. $\frac{2}{7}$

11. If $f(x) = x^2$, then domain of f is:

- a. Real number
- c. Rational number

- b. Integer
- d. Irrational

12. $2 \sin h x =$

- a. $e^x - e^{-x}$
- c. $\frac{ex - e^{-x}}{2}$

- b. $e^x + e^{-x}$
- d. $\frac{ex + e^{-x}}{2}$

13. $\lim_{\theta \rightarrow 0} \frac{\sin^2 \theta}{\theta} :$

- a. 0
- c. ∞

- b. 1
- d. 2

14. $x = at^2$, $y = 2at$ are the parametric equations of:

- a. Ellipse
- c. Parabola

- b. Circle
- d. Hyperbola

15. $\lim_{n \rightarrow \infty} \left(\frac{5n+1}{5n}\right)^n = :$

- a. $e^{1/5}$
- c. e^{-5}

- b. e^5
- d. $e^{-1/5}$

16. Let $f(x) = x^2 + \cos x$, then $f(x)$ is:

- a. Odd function
- c. Even function

- b. Constant function
- d. Neither even nor odd function

17. If at least one vertical line meets the curve at more than two points, then curve is:

- a. A function
- c. One-to-one function

- b. Not a function
- d. Onto function

CHAPTER : 2

- 1.** $\frac{d}{dx} \left(\frac{x^2 - 4}{x - 2} \right)$ equals to:
- a. 0
 - b. 1
 - c. $x+2$
 - d. $x-2$
- 2.** $\frac{d}{dx}(\tan x)$ is equals to:
- a. $\ln \cos x$
 - b. $-\ln \sin x$
 - c. $-\sec^2 x$
 - d. $\sec^2 x$
- 3.** The differential co-efficient of $e^{\sin x}$ equals:
- a. $e^{snx} \cdot \cos x$
 - b. $e^{snx} \cdot \sin x$
 - c. $e^{snx-1} \cdot \cos x$
 - d. $e^{snx-1} \cdot \sin x$
- 4.** If $y = \ln(\sin x)$, then $\frac{dy}{dx}$ equals:
- a. $\tan x$
 - b. $\cot x$
 - c. $-\tan x$
 - d. $-\cot x$
- 5.** If $f(x + h) = 2^{x+h}$, then $f'(x)$ equals to:
- a. $\frac{2^x}{\ln 2}$
 - b. 2^{x+h}
 - c. $2^x \cdot \ln 2$
 - d. 2^x
- 6.** $\cosh^{-1} x = :$
- a. $\ln(x + \sqrt{x^2 + 1})$
 - b. $\ln(x + \sqrt{x^2 - 1})$
 - c. $\frac{1}{2} \ln \left(\frac{1+x}{1-x} \right)$
 - d. $\frac{1}{2} \ln \left(\frac{x+1}{x-1} \right)$
- 7.** If $f(x) = x^{2/3}$ then $f'(8) = :$
- a. $\frac{1}{2}$
 - b. $\frac{2}{3}$
 - c. $\frac{1}{3}$
 - d. 3
- 8.** $\frac{d}{dx} (\cos x) = :$
- a. $\cos x$
 - b. $\sec^2 x$
 - c. $-\sin x$
 - d. $\sec^2 x$

- 9.** If $4y + 3x + 7 = 0$ then $\frac{dy}{dx}$ is :
- a. $\frac{3}{4}$
 - b. $-\frac{3}{4}$
 - c. $-\frac{4}{3}$
 - d. $\frac{2}{3}$
- 10.** If $y = \ln x$ then $\frac{dy}{dx} = :$
- a. $-\frac{1}{x}$
 - b. x^{-1}
 - c. x
 - d. $-2x$
- 11.** If $y = \sin 3x$ then $y_2 = :$
- a. $3\cos 3x$
 - b. $9\cos 3x$
 - c. $-9\sin 3x$
 - d. $9\sin 3x$
- 12.** If $y = x^3$ then $dy = :$
- a. $x^3 dx$
 - b. $3x^2 dx$
 - c. $\frac{x^4}{4} dx$
 - d. $x dx$
- 13.** $\lim_{\delta x \rightarrow 0} \frac{f(x+\delta x) - f(x)}{\delta x}$ is equal to:
- a. $f(0)$
 - b. $f(a)$
 - c. $f'(x)$
 - d. $f'(0)$
- 14.** $\frac{d}{dx} x^n$ is equal to:
- a. nx^{n-1}
 - b. x^{n-1}
 - c. $\frac{x^{n+1}}{n}$
 - d. nx^{n+1}
- 15.** If $f(x) = \cos x$ then $f'(0)$ is equal to:
- a. 0
 - b. -1
 - c. 1
 - d. $\frac{1}{2}$
- 16.** $\frac{d}{dx} \sin h^{-1} x$ is equal to:
- a. $\frac{1}{\sqrt{1-x^2}}$
 - b. $\frac{-1}{\sqrt{1-x^2}}$
 - c. $\frac{1}{\sqrt{1+x^2}}$
 - d. $\frac{-1}{\sqrt{1+x^2}}$
- 17.** If $f(x) = e^{ax}$ then $f'(x)$ is equal to:

a. $\frac{e^{ax}}{a}$
 c. $a e^{ax}$

b. $-\frac{e^{ax}}{a}$
 d. $-a e^{ax}$

18. $\frac{d}{dx} [\ln x]$ is equal to:

- a. X
 c. x^2

b. $\frac{1}{x}$
 d. $\frac{1}{x^2}$

19. Derivative of $\sin^{-1} x$ w.r.t x equals:

a. $\frac{1}{\sqrt{1-x^2}}$
 c. $\frac{1}{\sqrt{1+x^2}}$

b. $\frac{-1}{\sqrt{1-x^2}}$
 d. $\frac{-1}{\sqrt{1+x^2}}$

20. $\frac{d}{dx} (e^{\cos x})$ equals:

- a. $-\sin x e^{\cos x}$
 c. $\cos x e^{\sin x}$

b. $\sin x e^{\cos x}$
 d. $-\cos x e^{\sin x}$

21. If $f(x + h) = \cos(x+h)$, then $f'(x)$ equals:

- a. $\cos x$
 c. $\sin x$

b. $-\cos x$
 d. $-\sin x$

22. If $y = \sec\left(\frac{3\pi}{2} - x\right)$, then y' equals:

- a. cosec x cot x
 c. sec x tan x

b. -cosec x cot x
 d. -sec x tan x

23. $\frac{d}{dx} \tan h^{-1} x$ equals:

a. $\frac{1}{1+x^2}$
 c. $\frac{1}{1-x^2}$

b. $\frac{1}{x^2-1}$
 d. $\frac{1}{\sqrt{x^2-1}}$

24. The expression $\ln(x + \sqrt{x^2 + 1})$ equals:

- a. $\sin h^{-1} x$
 c. $\tan h^{-1} x$

b. $\cos h^{-1} x$
 d. cosec $h^{-1} x$

25. The differential co-efficient of $e^{\sin x}$ is:

- a. $e^{\sin x} \cdot \cos x$
 c. $e^{\cos x} \cdot \cos x$

b. $e^{\sin x} \cdot \sin x$
 d. $\sin x \cdot e^{\sin x-1}$

26. If $f(x) = 2^x$, then $f'(x) = :$

- a. 2^{x-1}
- b. $2^x \ln 2$
- c. $\frac{2^x}{\ln 2}$
- d. $\frac{\ln 2}{2^x}$

27. If $f'(c) = 0$, then $f(x)$ has relative maxima at $x = c$, if:

- a. $f'(c) < 0$
- b. $f'(c) = 0$
- c. $f'(c) > 0$
- d. $f'(c) \geq 0$

28. If $y = \ln(\sin x)$, then $\frac{dy}{dx}$ equals:

- a. $\tan x$
- b. $\cot x$
- c. $-\tan x$
- d. $-\cot x$

29. $\frac{d}{dx}(a^x)$ is:

- a. $\frac{\ln a}{a^x}$
- b. $\frac{a^x}{\ln a}$
- c. a^x
- d. $a^x \ln a$

30. If $y = \frac{1}{x^2}$, then $\frac{dy}{dx}$ at $x = -1$ is:

- a. 2
- b. 3
- c. $\frac{1}{3}$
- d. 4

31. $f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^n(0)}{n!} + \dots$ is called:

- a. Taylor series
- b. Binomial series
- c. Laurent series
- d. Maclaurin series

32. $\frac{1}{1+x^2}$ is derivative of:

- a. $\sin^{-1} x$
- b. $\sec^{-1} x$
- c. $\tan^{-1} x$
- d. $\cot^{-1} x$

33. $\frac{d}{dx}(\sec x)$ is:

- a. $\sec^2 x$
- b. $-\sec x x \tan x$
- c. $\sec x x \tan x$
- d. $\sec x \cot x$

34. $\frac{d}{dx} \cot^2 2x =:$

- a. $4 \cot 2x \operatorname{cosec} 2x$
- b. $-4 \cot 2x \operatorname{cosec}^2 2x$
- c. $4 \cot^2 2x \operatorname{cosec} 2x$
- d. $-4 \cot 2x$

35. $\frac{d}{dx} e^{x+h}$

- a. $\frac{e^{x+h}}{\ln h}$
c. e^{x+h}

- b. $\frac{e^{x+h}}{\ln x}$
d. he^{x+h}

36. If $f(x) = \cos x$ then $f'(\frac{\pi}{2}) = :$

- a. 1
c. $\frac{1}{2}$

- b. 0
d. -1

37. $\frac{d}{dx} (x^2 + 1)^2 = :$

- a. $1 - \frac{1}{2x}$
c. 0

- b. $1 + \frac{1}{2x}$
d. $1 - \frac{1}{x^2}$

38. $\frac{d}{dx} (\sqrt{x} - \frac{1}{\sqrt{x}})^2 = :$

- a. $2(x^2 + 1)$
c. $2x(x^2 + 1)$

- b. $\frac{(x^2+1)^3}{3}$
d. $4x(x^2 + 1)$

39. If $f(x) = \cos h x$ then $f(x)^2 - f'(x)^2 = :$

- a. 0
c. $\frac{1}{2}$

- b. 1
d. 2^2

40. $\frac{d}{dx} ((\ln x)^m)^k :$

- a. $\frac{mk}{x} (\ln x)^{mk-1}$
c. $\frac{1}{x^{mk}}$

- b. $\frac{k}{x^m} (\ln x)^{k-1}$
d. $\frac{mk}{x}$

41. $\frac{d}{dx} (\sqrt{x} - \frac{1}{\sqrt{x}})^2 :$

- a. $1 - \frac{1}{2x}$
c. 0

- b. $1 + \frac{1}{x^2}$
d. $1 - \frac{1}{x^2}$

42. If $y = \cos x$, $u = \sin x$ then $\frac{dy}{dx} = :$

- a. $\cos x$
c. $-\tan x$

- b. $-\cot x$
d. $-\operatorname{cosec} x$

- 43.** $a_0 + a_1 + a_2x^2 + \dots + a_nx^n + \dots$ is:
- a. Maclaurin's series
 - b. Taylor series
 - c. **Power series**
 - d. Binomial series
- 44.** $\frac{d}{dx} \sin^{-1} x = :$
- a. $\frac{1}{\sqrt{1+x^2}}$
 - b. $\cos^{-1} x$
 - c. $\frac{1}{\sqrt{1-x^2}}$
 - d. $\frac{1}{\sqrt{1-x}}$
- 45.** The order of the differential equation $\frac{d^2y}{dx^2} - \frac{dy}{dx} + 2x = 0$ is:
- a. 2
 - b. 1
 - c. 0
 - d. 3
- 46.** $\cos h^2 x - \sin h^2 x = :$
- a. 1
 - b. -1
 - c. 0
 - d. 2
- 47.** If $f(x)$ has second derivative at "c" such that $f'(c) = 0$ and $f''(c) < 0$ then "c" is a point of:
- a. **Maxima**
 - b. Minima
 - c. Zero point
 - d. Point of inflection
- 48.** If $y = \sqrt{1 - x^2}$, $0 < x < 1$ then $\frac{dy}{dx} = :$
- a. $\sqrt{x^2 - 1}$
 - b. $\frac{1}{\sqrt{x^2-1}}$
 - c. $\frac{x}{\sqrt{1-x^2}}$
 - d. $\frac{-x}{\sqrt{1-x^2}}$
- 49.** If $y = e^{\sin x}$, then $\frac{dy}{dx} = :$
- a. $e^{\sin x}$
 - b. $e^{\sin x} \cos x$
 - c. $e^{\sin x} + \cos x$
 - d. $-e^{\sin x} \cos x$
- 50.** $\frac{d}{dx} \cos h x = :$
- a. $-\sin hx$
 - b. $\sec hx$
 - c. $-\sec hx$
 - d. $\sin hx$
- 51.** Solution of $\frac{dy}{dx} = \frac{-y}{x}$ is:

a. $\frac{x}{y} = c$
c. $y = cx$

b. $\frac{y}{x} = c$
d. $xy = c$

52. If $x = f(\theta), y = g(\theta)$, then $\frac{dy}{dx}$:

a. $\frac{dy}{d\theta} \frac{d\theta}{dx}$
c. $\frac{d\theta}{dy} \frac{dx}{d\theta}$

b. $\frac{dx}{d\theta} \frac{d\theta}{dy}$
d. $\frac{dy}{d\theta} \frac{dx}{d\theta}$

53. $\frac{d}{dx} \log_a x =$

a. $\frac{1}{x}$
c. $\frac{1}{x} \ln a$

b. $x \ln x - x$
d. $\frac{1}{x \ln a}$

54. $\frac{d}{dx} \sec hx =:$

a. $\sec hx \tan h x$
c. $\tan h^2 x$

b. $-\sec hx \tan h x$
d. $\sec h^2 x$



CHAPTER : 3

1. $\int \frac{1}{x \ln x} dx$

- a. $\ln(\ln x) + c$
- c. $\ln\left(\frac{1}{2}\right) + c$
- b. $\ln x + c$
- d. $\ln(\ln \frac{1}{x}) + c$

2. Anti-derivative of $\cot x$ is equals to

- a. $\ln \cos x + c$
- c. $-\ln \cos x + c$
- b. $\ln \sin x + c$
- d. $-\ln \sin x + c$

3. $\int_0^3 \frac{1}{x^2+9} dx$ equals:

- a. $\frac{12}{\pi}$
- c. $-\frac{12}{\pi}$
- b. $\frac{\pi}{12}$
- d. $-\frac{\pi}{12}$

4. Solution of $y \cdot dx + x \cdot dy = 0$ is equal to:

- a. $x \cdot y = \text{constant}$
- c. $x+y = \text{constant}$
- b. $\frac{x}{y} = \text{constant}$
- d. $x-y = \text{constant}$

5. $\int (2x+3)^{1/2} dx$ is equal to:

- a. $\frac{1}{2} (2x+3)^{\frac{1}{2}} + c$
- c. $\frac{1}{3} (2x+3)^{\frac{1}{2}} + c$
- b. $\frac{2}{3} (2x+3)^{\frac{3}{2}} + c$
- d. $\frac{1}{3} (2x+3)^{\frac{3}{2}} + c$

6. $\int e^x \left(\frac{1}{x} + \ln x \right) dx$ equals:

- a. $e^{-x} \ln x + c$
- c. $e^x \cdot \frac{1}{x} + c$
- b. $e^{-x} \cdot \frac{1}{x} + c$
- d. $e^x \cdot \ln x + c$

7. For $n \neq -1$, $\int (f(x))^n f'(x) dx =$:

- a. $\frac{f'(x)}{n} + c$
- c. $\frac{(f(x))^{n+1}}{n+1} + c$
- b. $\frac{(f(x))^{n-1}}{n-1}$
- d. $(f(x))^{n+1} + c$

8. $\int \frac{e^{\tan^{-1} x}}{1+x^2} dx :$

- a. $e^{\sec x} + c$
- c. $e^{\cot^{-1} x} + c$
- b. $e^{\tan x} + c$
- d. $e^{\tan^{-1} x} + c$

9. $\int \frac{-1}{\sqrt{x^2-1}} dx =:$

- a. $\tan^{-1}x + c$
- c. $\sec^{-1}x + c$
- b. $\cosec^{-1}x + c$
- d. $\sin^{-1}x + c$

10. $\int \sec x dx =:$

- a. $\ln |\sec x + \tan x| + c$
- c. $\ln |\sec x - \tan x| + c$
- b. $\ln |\cosec x - \cot x| + c$
- d. $-\ln |\cosec x + \cot x| + c$

11. $\int_{-\pi}^{\pi} \sin x dx =:$

- a. 0
- c. 8
- b. 6
- d. 16

12. The integration is the reverse process of:

- a. Induction
- c. Tabulation
- b. Differentiation
- d. Sublimation

13. $\int \sin x dx$ is equal to:

- a. $\cos x$
- c. $-\sin x$
- b. $\sin x$
- d. $-\cos x$

14. $\int \frac{f'(x)}{f(x)} dx$ is equal to:

- a. $\ln x$
- c. $\ln f(x)$
- b. $\ln f(x)$
- d. $f(x)$

15. $\int \sec x \tan x dx$ is equal to:

- a. $\tan x$
- c. $\tan^2 x$
- b. $\sec^2 x$
- d. $\sec x$

16. $\int_0^1 x^3 dx$ is equal to:

- a. 4
- c. $\frac{1}{4}$
- b. -4
- d. $-\frac{1}{4}$

17. Solution of differential equation, $\frac{dy}{dx} = y$ is:

- a. ce^x
- c. e^x
- b. ce^{-x}
- d. e^{-x}

18. $\int \sin x \, dx$ is equal to:

- a. $\cos x$
- c. $\sin x$
- b. $-\cos x$
- d. $-\sin x$

19. $\int \frac{1}{1+x^2} \, dx$ is equal to:

- a. $\tan^{-1} x$
- c. $\cot^{-1} x$
- b. $\tan^{-1} x^2$
- d. $\cot^{-1} x^2$

20. $\int_a^b x \, dx$ equals:

- a. $\frac{b-a}{2}$
- c. $\frac{b^2-a^2}{2}$
- b. $\frac{b+a}{2}$
- d. $\frac{b^2+a^2}{2}$

21. $\int \frac{1}{x \ln x} \, dx$ equals:

- a. $\ln(\ln x)$
- c. $x \ln x$
- b. $\ln x$
- d. $\frac{\ln x}{x}$

22. $\int_1^4 \sqrt[3]{x} \, dx$ is equal to:

- a. 1
- c. 14
- b. 4
- d. 41

23. $\int e^{2x} (-\sin x + 2 \cos x) \, dx$ equals:

- a. $e^{2x} \sin x$
- c. $-e^{2x} \sin x$
- b. $e^{2x} \cos x$
- d. $-e^{2x} \cos x$

24. $\int_0^{\pi/4} \frac{\sec^2 x}{1+\tan x} \, dx$:

- a. 1
- c. $\ln 2$
- b. 2
- d. $\ln \sqrt{2}$

25. $\int (2x+3)^{1/2} \, dx$ equals:

- a. $\frac{1}{2}(2x+3)^{1/2} \, dx$
- c. $\frac{1}{3}(2x+3)^{1/2} + c$
- b. $\frac{2}{3}(2x+3)^{3/2} + c$
- d. $\frac{1}{3}(2x+3)^{3/2} + c$

26. The solution of differential equation $\frac{dy}{dx} = \sec^2 x$ is:

- a. $y = \cos x + c$
- c. $y = \cos^2 x + c$
- b. $y = \sec x + c$
- d. $y = \tan x + c$

27. Anti-derivative of $\cot x$, equals:

- a. $\ln(\cos x) + c$
- c. $-\operatorname{cosec}^2 x + c$
- b. $\ln(\sin x) + c$
- d. $\ln(\sec x) + c$

28. $\int_{-\pi}^{\pi} \sin x \, dx$ is equal to:

- a. 0
- c. 8
- b. 6
- d. 16

29. $\int \sin 3x \, dx$ is:

- a. $\frac{\cos 3x}{3} + c$
- c. $3 \cos 3x + c$
- b. $-\frac{\cos 3x}{3} + c$
- d. $-3 \cos 3x + c$

30. $\int_{-1}^3 x^3 \, dx$ is:

- a. 20
- c. 30
- b. 40
- d. 60

31. $\int \tan x \, dx$ is:

- a. $\ln \sec x + c$
- c. $\ln \sin x + c$
- b. $\ln \operatorname{cosec} x + c$
- d. $\ln \cot x + c$

32. $\int_a^b f(x) \, dx$ is:

- a. $-\int_a^b f(x) \, dx$
- c. $\int_{-b}^{-a} f(x) \, dx$
- b. $-\int_b^a f(x) \, dx$
- d. $\int_{-b}^{+a} f(x) \, dx$

33. $\int \frac{x}{x+2} \, dx = :$

- a. $\ln(x+2) + c$
- c. $x - 2 \ln(x+2) + c$
- b. $x+2 \ln(x+2) + c$
- d. $x - \ln(x+2) + c$

34. $\int e^x (\ln x + \frac{1}{x}) \, dx = :$

- a. $e^x \cdot \frac{1}{x} + c$
- c. $\frac{e^x}{\ln x} + c$
- b. $\frac{e^x}{\ln x} + c$
- d. $e^x \ln x + c$

35. $\int_0^1 \frac{1}{1+x^2} \, dx = :$

- a. 0
- c. $\frac{\pi}{4}$
- b. $\frac{\pi}{2}$
- d. $\frac{\pi}{3}$

- 36.** $\int \sec x \, dx =:$
- a. $\sec x \tan x + c$
 - b. $\sec^2 x \tan x + c$
 - c. $\ln(\sec x - \tan x + c)$
 - d. $\ln (\sec x + \tan x + c)$
- 37.** $\int \tan x \, dx =:$
- a. $\ln \cot x + c$
 - b. $\ln \cos x + c$
 - c. $\ln \sin x + c$
 - d. $\ln \sec x + c$
- 38.** $\int e^x \left[\frac{1}{1+x^2} + \tan^{-1} x \right] \, dx =:$
- a. $e^x \tan x + c$
 - b. $\frac{e^x}{1+x^2} + c$
 - c. $e^x \sin x + c$
 - d. $e^x \tan^{-1} x + c$
- 39.** $\int_0^{\pi/2} \sin^3 x \cos x \, dx =:$
- a. $\frac{1}{2}$
 - b. $\frac{2}{3}$
 - c. $\frac{1}{4}$
 - d. $\frac{1}{9}$
- 40.** The solution of $\frac{dy}{dx} = -y$ is:
- a. $y = e^2 x$
 - b. $y = ce^{-x}$
 - c. $y = e^x$
 - d. ce^x
- 41.** $\int \frac{1}{f(x)} \times f'(x) \, dx =:$
- a. $\ln x + c$
 - b. $\ln [f(x) + c]$
 - c. $\frac{1}{f(x)} + c$
 - d. $\ln |f(x)| + c$
- 42.** $\int 3^x \, dx =:$
- a. $3^x + c$
 - b. $3^x + \ln 3 + c$
 - c. $\frac{3^x}{\ln 3} + c$
 - d. $3 \ln 3^x + c$
- 43.** $\int_0^{\pi/2} \cos x \, dx =:$
- a. 0
 - b. 1
 - c. 2
 - d. 3
- 44.** $\int \sec^2 x \, dx:$
- a. $\tan x$
 - b. $\frac{\sec^3 x}{3}$
 - c. \tan^2
 - d. $\sec x \tan x$

45. Domain of $f(x) = x^2 + 1$:

- a. \mathbf{R}
- b. $\mathbf{R} - \{1\}$
- c. $\mathbf{R} - \{-1\}$
- d. $[1, \infty]$

46. $\int \sin x \cos x \, dx$:

- a. $\frac{1}{2} \cos 2x$
- b. $-\frac{1}{2} \cos 2x$
- c. $\frac{\sin^2 x}{2}$
- d. $\frac{\cos^2 x}{2}$

47. $\int \frac{1}{\sqrt{x^2-1}} \, dx$:

- a. $\sin^{-1} x$
- b. $\tan^{-1} x$
- c. $\sec^{-1} x$
- d. $\cosec^{-1} x$



CHAPTER : 4

1. Slope of line perpendicular to line $2x - 3y + 1 = 0$ is equal to:
- a. $\frac{3}{2}$
 - b. $-\frac{3}{2}$
 - c. $\frac{2}{3}$
 - d. $-\frac{2}{3}$
2. X- coordinate of centroid of triangle ABC with A (-2,3) ; B (-4,1); C (3,5) equals to:
- a. -1
 - b. 1
 - c. 3
 - d. -3
3. For any point (x,y) on x-axis:
- a. $y = 0$
 - b. $y = -1$
 - c. $y = 1$
 - d. $y = 2$
4. The point of concurrency of medians of triangle is called:
- a. In-centre
 - b. Centroid
 - c. Circumcentre
 - d. Orthocenter
5. The point of intersection of medians of a triangle is called:
- a. In-centre
 - b. Centroid
 - c. Circumcentre
 - d. Orthocenter
6. The distance of point P(1,6) from the line $6x - 4y + 9 = 0$ is:
- a. 49
 - b. $\frac{49}{52}$
 - c. $\frac{\sqrt{49}}{52}$
 - d. $\frac{49}{\sqrt{52}}$
7. Slope intercept form of line equals:
- a. $y - y_1 = m(x - x_1)$
 - b. $\frac{x}{a} + \frac{y}{b} = 1$
 - c. $x \cos \theta + y \sin \theta = p$
 - d. $y = mx + c$
8. Point of interception of lines $x - 2y + 1 = 0$ and $2x - y + 2 = 0$
- a. (1, 0)
 - b. (0, 1)
 - c. (-1, 0)
 - d. (0, -1)
9. Distance between (1,2) and (2,1) is:
- a. 1
 - b. 2
 - c. $\sqrt{2}$
 - d. 0

10. The line $y = mx + c$, will be tangent to the parabola $y^2 = 4ax$ if:

- a. $c = -am^2$
- b. $c = \frac{a}{m}$
- c. $c = a(1 + m^2)$
- d. $c = \frac{m}{a}$

11. Equation of the line parallel to $x + 3y - 9 = 0$:

- a. $3x - y - 9 = 0$
- b. $3x + 9y + 7 = 0$
- c. $2x - 6y - 18 = 0$
- d. $x - 3y + 9 = 0$

12. $ax + by + c = 0$, will represent equation of straight line parallel to y-axis if:

- a. $a = 0$
- b. $b = 0$
- c. $c = 0$
- d. $a = 0, c = 0$

13. If a straight line is parallel to y-axis, then its slope is:

- a. -1
- b. 0
- c. 1
- d. Undefined

14. A linear equation in two variables represents:

- a. Circle
- b. Ellipse
- c. Hyperbola
- d. Straight line

15. The centroid of the triangle whose vertices are $(3, -5)$, $(-7, 4)$ and $(10, -2)$ is:

- a. $(-2, -2)$
- b. $(-2, 2)$
- c. $(2, -1)$
- d. $(0, 0)$

16. Two lines represented by $ax^2 + 2hxy + by^2 = 0$ are parallel if:

- a. $h^2 - ab = 0$
- b. $h^2 + ab = 0$
- c. $a + b = 0$
- d. $a - b = 0$

17. If m_1 and m_2 are slopes of two lines then lines are perpendicular if:

- a. $m_1 m_2 = 1$
- b. $m_1 m_2 + 1 = 0$
- c. $m_1 m_2 - 1 = 0$
- d. $m_1 + m_2 = 0$

18. The distance between the points $(0, 0)$ and $(1, 2)$ is:

- a. 0
- b. 2
- c. $\sqrt{3}$
- d. $\sqrt{5}$

19. Two non-parallel lines intersect each other:

- a. 1 point
- b. 0 point
- c. ∞ points
- d. 2 points

20. The equation of horizontal line through (a,b) is:

- | | |
|------------|------------|
| a. $y = a$ | b. $y = b$ |
| c. $x = a$ | d. $x = b$ |

21. The two lines $a_1x + b_1y = c$; $a_2x + b_2y = c_2$ are parallel if:

- | | |
|--------------------------|--------------------------|
| a. $a_1 - a_2 = 0$ | b. $a_1 - b_1 = 0$ |
| c. $a_1b_1 - a_2b_2 = 0$ | d. $a_1b_2 - a_2b_1 = 0$ |

22. The inclination of $x = y$ is:

- | | |
|---------------|----------------|
| a. 30° | b. 60° |
| c. 45° | d. 180° |

23. If the line $(3x - y + 5) + k(2x - 3y - 4) = 0$ is parallel to y-axis, then $k =$:

- | | |
|-------------------|-------------------|
| a. $-\frac{1}{3}$ | b. $-\frac{1}{4}$ |
| c. $-\frac{1}{5}$ | d. 0 |

24. If α is the inclination of the line l , then $\frac{x - x_1}{\cos \alpha} = \frac{y - y_1}{\sin \alpha} = r$ (say) is called:

- | | |
|---------------------|-------------------|
| a. Point slope form | b. Normal form |
| c. Symmetric form | d. Intercept form |

25. The centroid of a triangle divides each median in ratio:

- | | |
|--------|--------|
| a. 2:1 | b. 1:2 |
| c. 2:3 | d. 1:1 |

26. The perpendicular distance of line $3x + 4y - 10 = 0$ from the origin is:

- | | |
|------------------|------|
| a. | b. 1 |
| c. $\frac{1}{2}$ | d. 2 |

27. The straight line $y = mx + c$ is tangent to the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ if:

- | | |
|--------------------------|--------------------------|
| a. $c^2 = a^2 m^2 - b^2$ | b. $c^2 = b^2 m^2 + a^2$ |
| c. $c^2 = b^2 m^2 - a^2$ | d. $c^2 = a^2 m^2 + b^2$ |

28. If α is the inclination of a line " l ", then it must be true that:

- | | |
|------------------------------------|--------------------------------------|
| a. $0 \leq \alpha < \frac{\pi}{2}$ | b. $\frac{\pi}{2} \leq \alpha < \pi$ |
| c. $0 \leq \alpha < \pi$ | d. $0 \leq \alpha < 2\pi$ |

29. Horizontal line through (7, -9) is:

- | | |
|------------|-------------|
| a. $x = 7$ | b. $x = -9$ |
| c. $y = 7$ | d. $y = -9$ |

30. Equation of line bisecting II and IV quadrant:

- | | |
|----------------------|----------------|
| a. $y = x$ | b. $y = -x$ |
| c. $y = \frac{1}{x}$ | d. $x + y = 1$ |

31. Joint equations of two lines is $ax^2 + 2hxy + by^2 = 0$, if θ is angle between them, then
 $\tan \theta =:$

- | | |
|-----------------------------------|-----------------------------------|
| a. $\frac{2\sqrt{h^2 + ab}}{a+b}$ | b. $\frac{2\sqrt{h^2 - ab}}{a+b}$ |
| c. $\frac{\sqrt{h^2 + ab}}{a+b}$ | d. $\frac{\sqrt{h^2 - ab}}{a+b}$ |

32. Set of all points equidistant from a fixed point form:

- | | |
|--------------|-------------|
| a. Ellipse | b. Parabola |
| c. Hyperbola | d. Circle |

33. Distance of (x_1, y_1) from line $ax + by + c = 0$ is:

- | | |
|---|---|
| a. $\frac{ax_1 + by_1 + c}{\sqrt{a^2 + b^2}}$ | b. $\frac{ax_1 + by_1 - c}{\sqrt{a^2 + b^2}}$ |
| c. $\frac{ax_1 + by_1 + c}{\sqrt{a+b}}$ | d. $\frac{ax_1 + by_1 - c}{\sqrt{a+b}}$ |



CHAPTER : 5

1. Point (1,2) lies in the region of inequality:

- a. $2x+y > 5$
- c. $2x+y < 3$



- b. $2x+y \geq 5$
- d. $2x+y < 3$

2. $x = 4$ is the solution of inequality:

- a. $x + 3 > 0$
- c. $-2x + 3 > 0$

- b. $x - 3 < 0$
- d. $x + 3 < 0$

3. (1, 0) is the solution of inequality:

- a. $7x + 2y < 8$
- c. $3x + 5y < 6$

- b. $x - 3y < 0$
- d. $-3x + 5y > 2$

4. A function which is to be maximized or minimized is called:

- a. Exponential function
- c. Quadratic function

- b. Linear function
- d. **Objective function**

5. Solution set of inequality $2x < 3$ is:

- a. $(-\infty, \frac{3}{2})$
- c. (∞, ∞)

- b. $(\frac{3}{2}, \infty)$
- d. $(-\frac{3}{2}, \frac{3}{2})$

6. A function which is to be maximized or minimized is called:

- a. Subjective function
- c. **Objective function**

- b. Quantitative function
- d. Qualitative function

7. (1,0) is solution of inequality:

- a. $9x + 2y < 8$
- c. $3x + 5y < 6$

- b. $-x + 3y < 0$
- d. $3x + 5y > 4$

8. $ax + b < c$ is :

- a. **Linear inequality**
- c. Equation

- b. Identity
- d. Not inequality

9. The feasible solution which maximizes or minimizes the objective function is called:

- a. Exact solution
- c. Final solution

- b. **Optimal solution**
- d. Objective solution

10. System of linear inequalities involved in the problem is called:

- a. Coefficients
- c. **Problem constraints**

- b. Solution
- d. Boundaries

CHAPTER : 6

1. The center of circle $(x+3)^2 + (y-2)^2 = 16$ equals:

- a. (-3,2)
- b. (3,-2)
- c. (3,2)
- d. (-3,-2)

2. The eccentricity of $\frac{y^2}{4} - x^2 = 1$ equals:

- a. $\frac{2}{\sqrt{5}}$
- b. $-\frac{2}{\sqrt{5}}$
- c. $\frac{\sqrt{5}}{2}$
- d. $-\frac{\sqrt{5}}{2}$

3. The radius of circle $x^2 + y^2 + 2gx + 2fy + c = 0$ is:

- a. $\sqrt{g^2 + f^2}$
- b. $\sqrt{g^2 - f^2 + c}$
- c. $\sqrt{g^2 + f^2 + c}$
- d. $\sqrt{g^2 + f^2} - c$

4. The vertex of parabola $(x-1)^2 = 8(y+2)$ is:

- a. (1, -2)
- b. (0, 1)
- c. (-1, -2)
- d. (1, 2)

5. The set of all points in the plane that are equally distant from a fixed point is called:

- a. Ellipse
- b. Parabola
- c. Hyperbola
- d. Circle

6. The parabola of $x^2 = y$ passes through a point:

- a. $\left(\frac{1}{2}, \frac{1}{2}\right)$
- b. $\left(\frac{1}{4}, \frac{1}{2}\right)$
- c. $\left(\frac{1}{2}, \frac{1}{4}\right)$
- d. $\left(\frac{1}{2}, -\frac{1}{2}\right)$

7. Equation of axis of a parabola $x^2 = 4ay$ is:

- a. $x = 0$
- b. $x = a$
- c. $y = 0$
- d. $y = a$

8. Length of tangent from (0,1) to $x^2 + y^2 + 6x - 3y + 3 = 0$ is called:

- a. 2
- b. 3
- c. 4
- d. 1

9. Length of latus rectum of ellipse $\frac{x^2}{16} + \frac{y^2}{9} = 1$ is:

- | | |
|---|---|
| a. $\frac{9}{2}$
c. $\frac{16}{9}$ | b. $\frac{9}{4}$
d. $\frac{9}{16}$ |
|---|---|

10. Centre of the circle $(x - 1)^2 + (y + 3)^2 = 3$ is:

- | | |
|--|--|
| a. (-1, -3)
c. (1, 3) | b. (-1, 3)
d. (1, -3) |
|--|--|

11. The eccentricity e of the hyperbola is:

- | | |
|--|--|
| a. $e = 0$
c. $e > 1$ | b. $e < 1$
d. $e = 1$ |
|--|--|

12. $x = a \sec \theta$, $y = b \tan \theta$ represents the parametric equations of:

- | | |
|---------------------------------------|---|
| a. Circle
c. Ellipse | b. Parabola
d. Hyperbola |
|---------------------------------------|---|

13. Length of each latus rectum of an ellipse is:

- | | |
|--|--|
| a. $\frac{2a^2}{b}$
c. $\frac{2b^2}{a}$ | b. $\frac{a^2}{b}$
d. $\frac{b^2}{a}$ |
|--|--|

14. If the ends of the diameter of the circle are (0,1) and (2,3), then its area is:

- | | |
|-------------------------------------|--------------------------------------|
| a. π
c. 4π | b. 2π
d. 8π |
|-------------------------------------|--------------------------------------|

15. The directrix of the parabola $x^2 = -8y$ is:

- | | |
|--|--|
| a. $x + 2 = 0$
c. $y + 2 = 0$ | b. $x - 2 = 0$
d. $y - 2 = 0$ |
|--|--|

16. The equation $x^2 + y^2 + 2gx + 2fy + c = 0$ represents a circle with centre:

- | | |
|--|---------------------------------------|
| a. (-g, -f)
c. (f, g) | b. (+g, -f)
d. (0,0) |
|--|---------------------------------------|

17. Axis of the parabola $x^2 = 4ay$ is:

- | | |
|--|--|
| a. $y = 0$
c. $x = y$ | b. $x = 0$
d. $x = 1$ |
|--|--|

18. For ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, (a > b)$, then eccentricity $e = :$

a. $\frac{\sqrt{a^2 - b^2}}{a}$

b. $\frac{\sqrt{a^2 + b^2}}{a}$

c. $\frac{\sqrt{b^2 - a^2}}{a}$

d. $\frac{\sqrt{b^2 - a^2}}{b}$

19. Focal chord perpendicular to axis parabola is called:

- a. Latus Rectum
c. Vertex

- b. Eccentricity
d. Axis



CHAPTER : 7

1. Work done by constant force \underline{F} during displacement \underline{d} is equal to:

- a. $\underline{F} \cdot \underline{d}$
- b. $\underline{F} \times \underline{d}$
- c. $\underline{r} \times \underline{F}$
- d. $\underline{F} \times \underline{r}$



2. $2\underline{i} \cdot (2\underline{j} \times \underline{k})$ equals:

- a. 0
- b. 2
- c. 4
- d. 6

3. Magnitude of vector $2\hat{i} + 3\hat{j} + 4\hat{k}$ is:

- a. 29
- b. $\sqrt{29}$
- c. 28
- d. $\sqrt{28}$

4. $\hat{i} \times \hat{i} =:$


- a. 1
- b. 2
- c. 0
- d. -1

5. A vector with magnitude 1 is called:

- a. Null vector
- b. Unit vector
- c. Zero vector
- d. Constant vector

6. $\underline{J} \cdot (\underline{k} \times \underline{i})$ is equal to:

- a. 0
- b. -1
- c. 1
- d. 2



7. Moment of Force vector \underline{F} about (\underline{r}) is:

- a. $\underline{r} \times \underline{F}$
- b. $\underline{F} \times \underline{r}$
- c. $\underline{r} \cdot \underline{F}$
- d. $\underline{F} \cdot \underline{r}$

8. $2\underline{i} \cdot (2\underline{j} \times \underline{k})$ equals:

- a. 4
- b. 3
- c. 2
- d. 1

9. $(\hat{i} \times \hat{k}) \times \hat{j}$ equals:

- a. -1
- b. 0
- c. 1
- d. ∞

10. If α, β, γ be the direction angles of a vector then $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma$ equals:

- a. -1
- b. 0
- c. 1
- d. 2

11. The angle between the vectors $2\bar{i} + 3\bar{j} + \bar{k}$ and $2\bar{i} - \bar{j} - \bar{k}$ is:

- a. 90°
- b. 45°
- c. 60°
- d. 30°

12. $\hat{j} \times \hat{k}$ is equal to :

- a. $-\hat{i}$
- b. \hat{i}
- c. 0
- d. 1

13. An angle in a semi-circle is:

- a. Right angle
- b. Obtuse angle
- c. Reflective angle
- d. 0°

14. $2\bar{i} \cdot \bar{j} \times 2\bar{k} =$:

- a. 0
- b. 2
- c. 1
- d. 4

15. If vectors $2\hat{i} + \hat{j} + \hat{k}$ and $\hat{i} - 4\hat{j} + \alpha\hat{k}$ are perpendicular, then $\alpha =$:

- a. 1
- b. 2
- c. 3
- d. 4

16. $[\underline{a} \underline{b} \underline{a}]$ is equal to:

- a. 1
- b. \underline{a}
- c. 0
- d. \underline{b}

17. The direction cosines of y- axis are:

- a. $(0, 1, 0)$
- b. $(1, 0, 0)$
- c. $(0, 0, 1)$
- d. $(0, 0, 0)$

18. Length of the vector $2\bar{i} - \bar{j} - 2\bar{k}$ is:

- a. 2
- b. 4
- c. 3
- d. 5

19. Projection of vector \vec{u} on vector \vec{v} is:

- a. $\frac{\vec{u} \cdot \vec{v}}{|\vec{v}|}$
- b. $\frac{\vec{u} \cdot \vec{v}}{|\vec{u}|}$
- c. $\frac{\vec{u} \times \vec{v}}{|\vec{v}|}$
- d. $\frac{\vec{u} \times \vec{v}}{|\vec{u}|}$

20. If \vec{v} is any vector, then vector of magnitude 5 opposite to \vec{v} is :

- a. $5\vec{v}$
- b. $-5\vec{v}$
- c. $5\frac{\vec{v}}{|\vec{v}|}$
- d. $-5\frac{\vec{v}}{|\vec{v}|}$